

FIFRA §2(ee) RECOMMENDATION
**FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF ARKANSAS, CALIFORNIA,
 COLORADO, CONNECTICUT, DELAWARE, DISTRICT OF COLUMBIA, FLORIDA,
 GEORGIA, HAWAII, IDAHO, ILLINOIS, INDIANA, IOWA, KANSAS, KENTUCKY,
 LOUISIANA, MAINE, MARYLAND, MASSACHUSETTS, MICHIGAN, MINNESOTA,
 MISSISSIPPI, MISSOURI, MONTANA, NEBRASKA, NEVADA, NEW HAMPSHIRE, NEW
 JERSEY, NEW MEXICO, NEW YORK, NORTH CAROLINA, NORTH DAKOTA, OHIO
 OKLAHOMA, PENNSYLVANIA, PUERTO RICO, RHODE ISLAND, SOUTH CAROLINA,
 SOUTH DAKOTA, TENNESSEE, UTAH, VERMONT, VIRGINIA, WASHINGTON, WEST
 VIRGINIA, WISCONSIN, WYOMING**

GROUP 21A ACARICIDE



EPA REG. NO. 81880-29-10163

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- All applicable directions, restrictions, and precautions on the EPA registered label are to be followed.
- This recommendation must be in the possession of the user at the time of pesticide application.
- This recommendation is made as permitted under FIFRA §2(ee) and has not been submitted to or approved by the EPA.

CROP	PEST	RATE	COMMENTS
Ornamental plants and Foliage plants grown in greenhouses and outdoors	Greenhouse whitefly (<i>Trialeurodes vaporariorum</i>)	6.4 – 9.6 oz per acre OR 0.15 - 0.22 oz per 1000 sq ft	Apply using sufficient water volume to obtain complete and uniform coverage of all plant leaf surfaces. Apply when mites first appear and before a threshold of five spider mites per leaf is reached. Apply higher rates for whiteflies.
	<ul style="list-style-type: none"> • Do not apply more than 34.14 oz of product per acre, per year. • Avoid drift to all other crops and non-target areas. 		
Greenhouse tomatoes	Greenhouse whitefly (<i>Trialeurodes vaporariorum</i>)	6.4 – 9.6 oz per acre OR 0.15 - 0.22 oz per 1000 sq ft	Apply using sufficient water volume to obtain complete and uniform coverage of all plant leaf surfaces. Apply when mites first appear and before a threshold of five spider mites per leaf is reached. Apply higher rates for whiteflies.
	<ul style="list-style-type: none"> • PHI = 2 days • Do not make more than 2 applications per crop per year. • Do not apply more than 12.8 oz of product per crop per season. • Allow a minimum of 30 days between sequential applications 		
Greenhouse cucumbers	Greenhouse whitefly (<i>Trialeurodes vaporariorum</i>)	6.4 – 9.6 oz per acre OR 0.15 - 0.22 oz per 1000 sq ft	Apply using sufficient water volume to obtain complete and uniform coverage of all plant leaf surfaces. Apply when mites first appear and before a threshold of five spider mites per leaf is reached. Apply higher rates for whiteflies.
	<ul style="list-style-type: none"> • PHI = 1 day • Do not make more than 2 applications per crop per year. • Do not apply more than 19.2 oz of product per crop per season. • Allow a minimum of 30 days between sequential applications. • Not for use on cucumbers grown in the field. 		

Sanmite® is a registered trademark of Nissan Chemical Corporation.

25-06-01

EPA Reg. No. 81880-29-10163
 EPA Est. No.

Gowan®

Produced For:
 Gowan Company, LLC
 P. O. Box 5569
 Yuma, AZ 85366-5569