



Supplemental Label

MOCCASIN™ MTZ Herbicide (EPA Reg. No. 70506-335)

FOR USE ONLY IN THE STATE OF FLORIDA

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read and follow all applicable directions, restrictions and precautions on the federally-registered label.

This label and the container label must be in the possession of the user at the time of pesticide application. Use of MOCCASIN MTZ according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for MOCCASIN MTZ.

TOMATOES

Established Tomato Post-Directed and Row Middle Applications

MOCCASIN MTZ may be applied post-directed and to row middle in established tomatoes.

Soil Texture	MOCCASIN MTZ (Pt./A)
COARSE (Loamy sand, sandy loam)	1.75 – 3.0
MEDIUM (Loam, silt loam, silt, sandy clay, sandy clay loam)	3.0 – 3.8
FINE (Silty clay, silty clay loam, clay, clay loam)	3.0 – 4.56

The tomato plants must be at least 4 inches tall or 5 to 6 leaf stage at the time of application. Avoid contacting tomato foliage with spray. Apply specified dosage in 20 or more gallons of water per acre as a directed spray.

Precautions

- MOCCASIN MTZ may damage transplants that have been weakened by any cause. To prevent damage, plant only healthy transplants.
- For row-middle applications where tomatoes are grown on sandy soils and where high soil moisture conditions can exist (e.g., low binding and high evaporation conditions, there is potential for crop injury. The risk of injury can be reduced by:
 - Incorporating MOCCASIN MTZ immediately following application
 - Applying MOCCASIN MTZ seven or more days before transplanting (but only after the beds have been formed)
 - Minimizing the application of MOCCASIN MTZ onto the plastic of the bed
 - Any combination of the above

Restrictions

- Do not make more than one application per season.
- Do not harvest within 90 days of application if the single application rate of this product is MORE than 3.0 pts/A per year.
- Do not harvest within 30 days of application if the single application rate of this product is LESS THAN OR EQUAL to 3.0 pts/A per year.
- Do not apply within 3 days after periods of cool, wet or cloudy weather, or crop injury will occur.
- Do not use hot caps on tomatoes within 7 days before or at any time after application.
- Do not apply by air.
- Do not apply to varieties or cultivars with unknown tolerance to metribuzin or s-metolachlor.