

SPINOSAD

GROUP

5

INSECTICIDE

# FIFRA Section 24(c) Special Local Need Label



Corteva Agriscience LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

## STATIC® Spinosad ME

EPA Reg. No. 62719-592

EPA SLN No. FL-240002

For Distribution and Use Only in the State of Florida

**For Use Only By Division of Plant Industry, in Urban/Suburban/Agricultural Areas, in Proximity to Crops on Inanimate Objects (Such as Telephone Poles, Light Poles), Tree Trunks and Limbs to Eradicate Fruit Flies in the Family Tephritidae.**

### ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read this SLN and the label affixed to the container for this product before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of this product according to this FIFRA 24(c) Special Local Need labeling is subject to all use precautions and limitations imposed by the label affixed to the container for this product.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

#### Active Ingredients:

spinosad (a mixture of spinosyn A and spinosyn D) .....	2%
methyl eugenol.....	51%
Other ingredients: includes water, waxes and oils.....	47%
Total.....	100%

### Directions for Use

This SLN is intended for use as a control/eradication treatment only by federal or state personnel.

#### Dosage:

Apply 0.17-0.34 fl. oz. (5-10 ml.) of product per application site or dollops. Apply up to 600-1200 applications (spot treatments) per square mile. Do not apply more than 100 fl. oz. (2957 ml.) of product per acre per year (0.135 lb. a.i. spinosad), nor make more than 52 total applications per acre per year.

Bait Station - Apply 5-10 ml. of product per bait station.

**Dilution Rate:** Ready to use formula. Thoroughly stir product before each use. Do not dilute or mix with other materials.

**Method of Application:**

Spot Application by hand (spatula or other spreading equipment) or spot application using mechanical/pneumatic applicator that delivers large spot treatments.

**Frequency/Timing of Application:**

May be applied at 1-6 week intervals, depending on pest population monitoring and pest risk.

**Specific Use Restrictions:**

- Do not aerial apply this product.
- Do not apply this product to blooming trees or plants in order to protect bees.
- Do not apply STATIC® Spinosad ME to food or feed commodities.
- Do not apply to edible plants.
- All applications must be placed out of the reach of children and pets.
- Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark.
- Do not use in mixture with other pesticides unless provided for in the labeling.

**Expiration date:** 12/31/2029

™®Trademarks of Corteva Agriscience and its affiliated companies

R894-005

Approved: 03/21/2024

Replaces R894-003 and R894-004