



Recommendation

FIFRA SECTION 2(ee) RECOMMENDATION

GROUP	28	INSECTICIDE
-------	----	-------------

Shenzi 400 SC Insecticide

(EPA Reg. No. 70506-607)

FOR SUPPRESSION OF LESSER CORNSTALK BORER (*Elsamopalous lignosellus*) IN PEANUT IN THE STATES OF ALABAMA, ARKANSAS, GEORGIA, MISSISSIPPI, NORTH CAROLINA, SOUTH CAROLINA, FLORIDA, OKLAHOMA AND VIRGINIA

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by USEPA.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read and follow all applicable instructions, restrictions, and precautions, and Conditions of Sale and Warranty on the EPA-approved label. This label and the federally registered label must be in the possession of the user at the time of pesticide application.

CROP	APPLICATION METHOD	INSECT	SHENZI 400 SC INSECTICIDE RATE PER ACRE IN FL OZ (LB A.I.)	RE-TREATMENT INTERVAL IN DAYS (RTI)	PRE-HARVEST INTERVAL IN DAYS (PHI)	RE-ENTRY INTERVAL IN HOURS (REI)
Peanut	Foliar	Lesser Cornstalk Borer (<i>Elsamopalous lignosellus</i>)	1.7 - 3.8 (0.044 - 0.098)	5	1	4

APPLICATION INSTRUCTIONS:

Apply higher rates within the listed range for heavier infestations, larger/denser crops, or extreme environmental conditions such as rainy weather and high temperatures.

RESTRICTIONS:

- DO NOT make more than 4 applications per acre per calendar year.
- DO NOT apply more than 7.7 fl oz of SHENZI SC INSECTICIDE or 0.2 lb a.i. chlorantraniliprole containing products per acre per calendar year.

DO NOT make more than two sequential applications of SHENZI SC INSECTICIDE before rotating to another registered insecticide having a different mode-of-action.