



2(ee) Recommendation

EPA Reg. No. 60063-81

FOR CONTROL OF TAR SPOT ON CORN (FIELD CORN, CORN GROWN FOR SEED, POPCORN)

For use in the States of Illinois, Indiana, Iowa, Kansas, Michigan, Minnesota, Missouri, Nebraska, North Dakota, Ohio, South Dakota, and Wisconsin.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ AND FOLLOW THIS SECTION 2(ee) AND ENTIRE ANDIAMO 230 FUNGIDE LABEL BEFORE USING.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

FAILURE TO FOLLOW THE USE DIRECTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN PLANT INJURY OR POOR DISEASE CONTROL.

DISEASES CONTROLLED	RATE PER ACRE	APPLICATION DIRECTIONS
Tar Spot (<i>Phyllachora maydis</i>)	4.0-6.0 fl. oz./A (0.06 – 0.09 lbs. a.i./A)	<p>This product may be applied early (V4-V8) or late (Vt-R3) prior to disease development. Apply as a curative application when disease incident does not exceed 5% of the plants.</p> <p>Apply as a foliar spray (a minimum of 10 GPA by ground or a minimum of 2 GPA by air) or via chemigation in sufficient water to obtain thorough coverage of plants. See mixing instructions restrictions below.</p> <p>A compatibility agent, another fungicide, or an insecticide approved for use on corn may be included, if needed. The use directions and restrictions of the added product must be followed. Always follow the more restrictive label.</p>

RESTRICTIONS

- **DO NOT** apply more than 6 fl. oz. of this product per acre per year.
- Maximum total amount of tetraconazole active ingredient (lbs. a.i./A) which may be applied from all products per acre per year: 0.09 lbs. a.i./A.
- **DO NOT** make more than one (1) application per year.
- **DO NOT** apply this product after corn growth stage R3 (brown silk/milk).
- **DO NOT** use adjuvants in sprays made between V8 (8 leaf collar) and VT (lowest branch of the tassel visible but silks have not emerged) growth stage.
- **Mixing Instructions: DO NOT** allow the ratio of water:product in field to exceed 1:1. Fill the spray tank (or chemigation nurse tank) half full with water before adding any product. **DO NOT** add product to the spray tank before adding water, this may result in a hazardous chemical reaction.
- **Restricted entry interval (REI):** 12 hours for all activities with the exception of 3 days for detasseling corn grown for seed.

NOTICE

The directions, restrictions and precautions on this and the entire Andiamo 230 Fungicide label must be read and followed carefully. This supplemental bulletin must be in the possession of the user at the time of application. Read the entire label including the **WARRANTY AND LIMITATION OF DAMAGES** prior to use.

2023-12-05 Andiamo 230 Fungicide 2(ee)
Expiration Date: 2029-12-31