

RESTRICTED USE PESTICIDE

Due to Acute Toxicity And Toxicity to Birds and Mammals.
For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

Product Bulletin



Corteva Agriscience LLC 9330 Zionsville Road Indianapolis, IN 46268-1054 USA 800-258-3033

Vydate C-LV

EPA Reg. No. 352-532

2(ee) Recommendation†

For Distribution and Use Only in the State of Idaho

Suppression of *Globodera pallida* (Pale Cyst Nematode) in Potatoes

ATTENTION

† This recommendation is made as permitted under FIFRA 2(ee) and has not been submitted to or approved by the EPA.

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- Read and follow all applicable directions for use, precautions, and limitations on the Vydate C-LV product label attached to the container for this product.

Directions for Use

For maximum crop protection, use a pre-plant fumigant, shanked-in, and follow the recommended Vydate C-LV treatment program.

Note: For best results, make all applications, other than in-furrow, via chemigation.

| |
|--|
| Start with a fumigant that is applied pre-plant using a soil injection (shank) system. |
| 34 to 68 fluid oz/acre in-furrow at planting |
| 34 fluid oz/acre at crop emergence |
| 34 fluid oz/acre at 1440 degree days F (800 DD C) |
| 34 fluid oz/acre 14 days later |
| Continue applying every 14 days until 7 days before digging. |

®™ Trademarks of Corteva Agriscience and its affiliated companies

R604-005

Issued: 06/10/2025

Initial Printing