



Product Name: Tandem™

Tandem™ is a combination of the following products:

Tandem™ A Herbicide

Tandem™ B Herbicide

Water Conditioner

The specimen labels are attached

COMPANY IDENTIFICATION:

Corteva Agriscience Canada Company

Suite 240, 115 Quarry Park Rd. SE

Calgary, AB

T2C 5G9

Canada

Customer Information Number: 800-667-3852

E-mail address: solutions@corteva.com

EMERGENCY TELEPHONE

24-Hour Emergency Contact: 1-888-226-8832

Local Emergency Contact: 1-888-226-8832

™ Trademarks of Corteva Agriscience and its affiliated companies



Tandem™ A

Herbicide

| | | |
|-------|---|-----------|
| GROUP | 2 | HERBICIDE |
|-------|---|-----------|

FOR SALE FOR USE IN THE PRAIRIE PROVINCES AND INTERIOR OF BRITISH COLUMBIA (INCLUDING THE PEACE RIVER REGION) ONLY

AGRICULTURAL

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN

ACTIVE INGREDIENT: pyroxsulam 30 g/L
Suspension
Warning, contains the allergen soy

REGISTRATION NO. 29985 PEST CONTROL PRODUCTS ACT

WARNING



POISON

**EYE AND SKIN IRRITANT
POTENTIAL DERMAL SENSITIZER**

NET CONTENTS: 8 L - bulk

Corteva Agriscience Canada Company
Suite 240, 115 Quarry Park Rd. SE
Calgary, AB
T2C 5G9
1-800-667-3852

™Trademark of Corteva Agriscience and its affiliated companies.

PRECAUTIONS

HAZARD TO HUMANS AND DOMESTIC ANIMALS

KEEP OUT OF REACH OF CHILDREN

EYE AND SKIN IRRITANT

POTENTIAL DERMAL SENSITIZER

Harmful if swallowed. Causes eye and skin irritation. Harmful if inhaled. Do not get in eyes or on skin. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

At all times: Wear clean clothing with full length sleeves and pants.

Wear coveralls over long sleeved shirt and long pants, chemical-resistant gloves, socks and chemical resistant footwear during mixing, loading, application, clean-up and repair. Wear goggles or face shield during mixing/loading. For closed cab or aerial application, coveralls and gloves are not necessary.

Apply only when potential for drift to areas of human habitation and human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversion, application equipment and sprayer settings.

At completion of spraying or end of the day: Take a shower immediately. Wash thoroughly with soap and water before eating, drinking or smoking. Remove contaminated clothing at the end of the work session and store and wash separately from household laundry using detergents and hot water before reuse.

FIRST AID

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give **any** liquid to the person. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

TOXICOLOGICAL INFORMATION

No specific antidote. Employ supportive care. Treatment should be based on judgment of the physician in response to reactions of the patient. This product contains petroleum distillates. Vomiting may cause aspiration pneumonia.

AGRICULTURAL CHEMICAL

Do not ship or store with food, feeds, drugs or clothing.

ENVIRONMENTAL PRECAUTIONS

TOXIC to aquatic organisms and non-target terrestrial plants. Observe spray buffer zones specified under DIRECTIONS FOR USE. To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay. Avoid application when heavy rain is forecast. Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body. This product demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

STORAGE

Store in original container in a secure, dry, heated storage. This product will freeze at 14°F/-10°C. Do not store below 15°F/-9°C. Allow product to warm above 45°F/7°C before using and thoroughly mix the product prior to use. Do not allow contamination of seeds, plants, fertilizers or other pesticides. Do not contaminate food, feedstuffs or domestic water supplies. To prevent contamination, store this product away from food or feed.

DISPOSAL

Recyclable Containers

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial/territorial requirements.

Refillable Containers

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

Returnable Containers

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial/territorial regulatory agency. Contact the manufacturer and the provincial/territorial regulatory agency in case of a spill, and for clean-up of spills.

GENERAL INFORMATION

Tandem A Herbicide is a selective postemergence herbicide for the control of wild oats and certain broadleaved weeds in spring wheat, durum wheat and winter wheat not underseeded with legumes. Tandem A Herbicide is mixed with water and applied as a uniform broadcast spray. It is non-corrosive, nonflammable, and nonvolatile.

Tandem A Herbicide must be applied early postemergence, to the main flush of actively growing wild oats and broadleaved weeds. Conditions favouring active weed growth enhance the activity of Tandem A Herbicide by allowing maximum foliar uptake and contact activity. Weeds hardened off by cold weather or drought stress may not be adequately controlled or suppressed and re-growth may occur. For best results, ensure thorough spray coverage of target weeds. See "DIRECTIONS FOR USE" section of this label for complete use details.

Tandem A Herbicide stops growth of susceptible weeds rapidly. However, typical symptoms (discolouration) of dying weeds may not be noticeable for 1 to 3 weeks after application, depending upon growing conditions and weed susceptibility. Degree of control and duration of effect are dependent on weed sensitivity, weed size, crop competition, growing conditions at and following treatment, and spray coverage.

MODE OF ACTION

Tandem A Herbicide inhibits the production of the ALS enzyme in plants. This enzyme is essential for the production of certain amino acids which are essential for plant growth. Tandem A Herbicide is a Group 2 mode of action herbicide.

GENERAL USE PRECAUTIONS

Non-Target Sites

Do not apply where proximity of susceptible crops (e.g. canola and legumes) or other desirable plants is likely to result in exposure to spray or spray drift. See Environmental Precautions section of the label.

Crop Rotation

Fields previously treated with Tandem A Herbicide can be seeded 11 months following an application of Tandem A Herbicide, with the following crops: barley, brown mustard, canola, dry bean (of the species *Phaseolus vulgaris*), flax, canola quality Brassica juncea, lentils, oats, field peas, chickpea, spring wheat, soybean and yellow mustard or fields can be summerfallowed. Fields previously treated with Tandem A can be seeded 10 months following an application of Tandem A with the following crops: sunflower and potatoes (not for seed).

Tank Mixtures

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact Corteva Agriscience Canada Company at 1-800-667-3852 or www.corteva.ca for information before applying any tank mix that is not specifically recommended on this label.

Spray Equipment Precaution

Do not apply through any type of irrigation system.

Sprayer Clean-Out Instructions

To avoid injury to desirable plants, thoroughly clean equipment used to apply this product before re-use or using it to apply other chemicals.

1. Immediately after spraying, completely drain the sprayer tank. Any contamination on the outside of the spraying equipment should be removed by washing with clean water.
2. First rinse:
 - Spray the inside of tank with clean water and fill the sprayer with at least one tenth of the spray tank volume.
 - Agitate and circulate for 15 minutes, and flush through booms and hoses.
 - Remove end caps or open ball valves on the ends of each boom section, and flush solution through the boom ends to ensure there is no spray solution trapped between the boom end and the nozzles.
 - Drain tank completely.
3. Second rinse:
 - Fill the tank with clean water.
 - Add All Clear Spray Tank Decontaminator as per manufacturer's recommendations while filling the tank with clean water.
 - Agitate and then flush the boom and hoses with the cleaning solution. Top up with water making sure the tank is completely full. Allow to stand for 15 minutes with agitation. Flush the solution out of the spray tank through the spray booms. Remove end caps or open ball valves on the ends of each boom section, and flush solution through the boom ends to ensure there is no spray solution trapped between the boom end and the nozzles.
 - If possible, let the solution stand in the sprayer tank and booms for an extended period of time, overnight if possible.
 - After flushing the boom and hoses, drain tank completely.

- Remove nozzles and screens and clean separately with a cleaning agent or an ammonia solution (100 mL in 10 L water).
4. Third rinse:
- Rinse the tank with clean water and flush through the boom and hoses using at least one tenth of the spray tank volume.
 - Remove end caps or open ball valves on the ends of each boom section, and flush solution through the boom ends to ensure there is no spray solution trapped between the boom end and the nozzles.
 - Drain tank completely.

DIRECTIONS FOR USE

FOR SALE FOR USE IN THE PRAIRIE PROVINCES AND PEACE RIVER REGION OF BRITISH COLUMBIA ONLY

READ THE ENTIRE LABEL BEFORE USE. FAILURE TO FOLLOW LABEL INSTRUCTIONS MAY RESULT IN ERRATIC WEED CONTROL OR CROP DAMAGE. DO NOT APPLY TO CROPS UNDERSEED WITH LEGUMES.

As this product is not registered for the control of pests in aquatic systems, **DO NOT** use to control aquatic pests.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

Field sprayer application

DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) coarse classification. Boom height must be 60 cm or less above the crop or ground.

Aerial application

DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply when wind speed is greater than 16 km/h at flying height at the site of application. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) coarse classification. Reduce drift caused by turbulent wingtip vortices. Nozzle distribution along the spray boom length **MUST NOT** exceed 65% of the wing- or rotorspan.

TANDEM A APPLIED ALONE

Application Directions - Spring Wheat, Durum wheat

Apply Tandem A Herbicide at a rate of 500 mL per hectare. Add non-ionic surfactant (Agral 90, AgSurf or Surf 92) at 0.25 v/v (0.25 L per 100 L of spray solution). Apply using aerial equipment or ground equipment that will assure uniform coverage. See table below for weeds species controlled. Tandem A Herbicide will control only emerged weeds. Weeds that germinate after application will not be controlled.

Weeds Controlled or Suppressed with Tandem A Herbicide

Weeds Controlled:

| | |
|---|--|
| barnyard grass (1-5 leaf) | hemp nettle (1-8 leaf) |
| brome, Japanese (1-6 leaf) | oats, wild (up to 4 leaf, 2 tiller) |
| canola, volunteer† (1-6 leaf) | pigweed, red-root (1-8 leaf) |
| chickweed, common (up to 10 cm) | round-leaved mallow (up to 6 leaf stage, <10 cm in size) |
| cleavers (up to 6 whorl) | smartweed (lady's-thumb, 1-5 leaf) |
| corn spurry (up to 2 whorl stage, <10 cm in height) | shepherd's purse (up to 30 cm tall) |
| cow cockle (up to 8 leaf) | stinkweed (up to 30 cm tall) |
| flixweed (up to 10 cm) | |

foxtail, yellow (1-5 leaf)
†will not control volunteer Imidazolinone-tolerant canola

Weeds Suppressed:

| | |
|--|---|
| brome, downy (2-6 leaf, 4 tillers) | foxtail, green (1-5 leaf) |
| dandelion (seedlings and over-wintered rosettes less than or equal to 20 cm) | wild buckwheat (1-4 leaf) |
| | Canada thistle (up to 30 cm tall, prebud) |
| | Russian thistle (up to 10 cm tall) |

Application Timing

Apply to actively growing wheat from the 3 leaf stage to before the flag leaf stage. Apply when weeds are actively growing: grass weeds up to the 4 leaf, 2 tiller stage and broadleaved weeds as indicated in the table above. Only weeds emerged at the time of treatment will be controlled. Best results are obtained from applications made to seedling weeds. Extreme growing conditions such as drought or near freezing temperature prior to, at or following time of application may reduce weed control and increase the risk of crop injury at all stages of growth. Under conditions of low crop and high weed density, control may be reduced.

Application Directions - Winter Wheat (Fall or Spring Application) for Downy Brome

Tandem A Herbicide applied in the fall will provide control of downy brome while a spring application will provide suppression. Apply Tandem A Herbicide at a rate of 500 mL per hectare. Add BASF Assist Oil Concentrate at 0.8% v/v (0.8 L/100 L of spray mixture) or Merge Adjuvant at 0.5% v/v (0.5 L/100 L of spray mixture; spring application only). Apply using aerial equipment or ground equipment that will assure uniform coverage. In addition to suppression of downy brome when applied in the spring Tandem A Herbicide will also control/suppress weeds listed in Table. Tandem A Herbicide will control only emerged weeds. Weeds that germinate after application will not be controlled.

Application Timing

Apply in the fall when winter wheat is in the 1-3 leaf stage or in the spring when winter wheat is in the 2-7 leaf, 4 tiller stage. For fall applications, apply when downy brome is actively growing at the 1 to 3 leaf stage; for spring applications, apply when weeds are actively growing at the stages indicated in Table. Only weeds emerged at the time of treatment will be controlled. Best results are obtained from applications made to seedling weeds. Extreme growing conditions such as drought or near freezing temperature prior to, at or following time of application may reduce weed control and increase the risk of crop injury at all stages of growth. Under conditions of low crop and high weed density, control may be reduced.

Occasionally slight yellowing or height reduction may be observed in the treated crop. These transient symptoms disappear within 14 days with no reduction to yield. Do not apply to crops suffering from drought, nutrient deficiency or exposed to frost or other agronomic factors affecting plant growth.

If foliage is wet at the time of application control may be decreased. Applications of Tandem A Herbicide are rainfast within 2 hours after application.

Mixing Instructions

1. Fill sprayer tank 1/2 full of water.
2. Start sprayer tank agitation.
3. Add the required amount of Tandem A Herbicide
4. Fill the sprayer tank with sufficient water to spray 50-100 L of spray mixture per hectare.
5. Add non-ionic surfactant.
6. Follow sprayer directions and precautions as outlined above, especially when applying next to sensitive crops (e.g. canola and legumes).
7. Follow sprayer clean-up directions.

Pre-harvest/Grazing Intervals

Livestock may be grazed on treated crops 7 days following application. Do not harvest the treated crop within 60 days after application.

TANK MIX COMBINATIONS – FOR USE IN SPRING WHEAT AND DURUM WHEAT ONLY

Tandem A Herbicide can be tank mixed with herbicides and/or fungicides as indicated in the table below.

Do not use non-ionic surfactant with Tandem A Herbicide when tank mixing with registered broadleaved herbicide or fungicide tank-mix partners.

Tank-Mix Combinations with Tandem A Herbicide at 500 mL/ha in Spring Wheat and Durum Wheat

| Herbicide Tank-Mix Partner | Fungicide Tank-Mix Partner | Rate of herbicide TM partner (Product /ha) | Additional Pests Controlled |
|---|-----------------------------------|--|--|
| Tandem A Herbicide alone | Tilt 250 E or Stratego 250 EC | 500 mL/ha Consult fungicide tank-mix partner label for fungicide rates. | consult fungicide tank-mix partner label for diseases controlled |
| Tandem B Herbicide | | 310 mL/ha | consult label |
| Exhilarate A Herbicide + MCPA Ester 600 + Tandem B Herbicide | | 25 g/ha 467 mL/ha 310 mL/ha | consult label |
| Curtail™ M Herbicide | | 1.5 L/ha | annual sowthistle (1-5 leaf) consult label |
| MCPA LV500 | | 0.7-1.1 L/ha | stork's-bill (1-8 leaf) †† wild buckwheat†+ weeds labelled for this rate range of MCPA |
| 2,4-D Ester | | 280 g ae/ha (e.g., 425 mL/ha of 2,4-D ester 700) | volunteer canola (1-6 leaf stage)* lamb's quarters (1-6 leaf stage) *Including Imidazolinone-tolerant volunteer canola |
| MCPA Ester | | 280 g ae/ha (e.g., 465 mL/ha of MCPA ester 600) | volunteer canola (1-6 leaf stage)* lamb's quarters (1-6 leaf stage) *Including Imidazolinone-tolerant volunteer canola |

†for larger weeds or heavy infestations use MCPA at 1.1 L/ha

††Use the 1.1 L/ha rate for control of stork's-bill

Tank-Mix Combinations with Tandem A Herbicide at 375 mL/ha* in Spring Wheat and Durum Wheat

| Herbicide Tank-Mix Partner | Rate of herbicide TM partner (Product /ha) | Additional Pests Controlled |
|---|---|------------------------------------|
| Tandem B Herbicide | 210 - 310 mL/ha | consult label |
| Exhilarate A Herbicide + MCPA Ester 600 + Tandem B Herbicide | 25 g/ha 467 mL/ha 210 mL/ha | consult label |

*Under Low Wild Oat Populations (<75 plants/m²)

Mixing Instructions

1. Fill sprayer tank 1/2 full of water.
2. Start sprayer tank agitation.
3. Add the required amount of Tandem A Herbicide
4. Add the correct amount of tank-mix partner and continue to agitate.
5. Fill the sprayer tank with the remaining amount of water required.

Notes: Add only the adjuvant recommended. Follow tank-mix partner label for order of mixing.

APPLICATION INSTRUCTIONS (GROUND AND AERIAL APPLICATIONS)

(1) Ground Application (Tandem A Herbicide applied alone and with all tank-mix partners)

Using ground equipment, apply Tandem A alone or with tank-mix partners as a broadcast treatment. Apply Tandem A Herbicide alone at the recommended rates in a spray volume of 50-100 L water / ha.

(2) Aerial Application

Directions for Use

Do not use human flaggers.

Use Tandem A Herbicide alone or in tank-mix with MCPA Ester as a broadcast treatment by air. Apply the recommended rates of Tandem A Herbicide and MCPA Ester as indicated above. Apply in a minimum spray volume of 30 L water / ha.

Apply only by fixed-wing or rotary aircraft equipment which has been functionally and operationally calibrated for the atmospheric conditions of the area and the application rates and conditions of this label.

Label rates, conditions and precautions are product specific. Read and understand the entire label before opening this product. Apply only at the rate recommended for aerial application on this label. Where no rate for aerial application appears for the specific use, this product cannot be applied by any type of aerial equipment.

Ensure uniform application. To avoid streaked, uneven or overlapped application, use appropriate swath marking devices.

Use Precautions

Apply only when meteorological conditions at the treatment site allow for complete and even coverage. Apply only under conditions of good practice specific to aerial application as outlined in the National Aerial Pesticide Application Manual, developed by the Federal/Provincial/Territorial Committee on Pest Management and Pesticides.

Operator Precautions

Do not allow the pilot to mix chemicals to be loaded onto the aircraft. Loading of premixed chemicals with a closed system is permitted.

It is desirable that the pilot have communication capabilities at each treatment site at the time of application.

The field crew and the mixer/loaders must wear chemical resistant gloves, coveralls and goggles or face shield during mixing/loading, cleanup and repair. Follow the more stringent label precautions in cases where the operator precautions exceed the generic label recommendations on the existing ground boom label.

All personnel on the job site must wash hands and face thoroughly before eating and drinking. Protective clothing should be laundered regularly.

Product Specific Precautions

Read and understand the entire label before opening this product. If you have questions, call the manufacturer at 1-800-667-3852 or obtain technical advice from the distributor or your provincial agricultural representative. Application of this specific product must meet and/or conform to the following:

- The interaction of many equipment-and-weather-related factors determine the potential for spray drift. Users are responsible for considering all these factors when making decisions.
- The following drift management requirement must be followed to avoid off-target drift movement from aerial applications:
Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.

SPRAY BUFFER ZONES

A spray buffer zone is NOT required for:

- uses with hand-held application equipment permitted on this label.

The spray buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands), and sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands).

| Method of application | Crop | | Spray Buffer Zones (metres) Required for the Protection of: | |
|-----------------------|---|-------------|---|---------------------|
| | | | Freshwater Habitat | Terrestrial Habitat |
| Field sprayer | Spring wheat, durum wheat, winter wheat | | 1 | 2 |
| Aerial | Spring wheat, durum wheat, winter wheat | Fixed wing | 1 | 65 |
| | | Rotary wing | 1 | 55 |

When tank mixes are permitted, consult the labels of the tank-mix partners and observe the largest (most restrictive) spray buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.

The spray buffer zones for this product can be modified based on weather conditions and spray equipment configuration by accessing the Spray Buffer Zone Calculator on the Pesticides portion of the Canada.ca website.

RESISTANCE MANAGEMENT RECOMMENDATIONS

For resistance management, Tandem A Herbicide is a Group 2 herbicide. Any weed population may contain or develop plants naturally resistant to Tandem A Herbicide and other Group 2 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance management strategies should be followed.

To delay herbicide resistance:

- Where possible, rotate the use of Tandem A Herbicide or other Group 2 herbicides within a growing season (sequence) or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group when such use is permitted. To delay resistance, the less resistance-prone partner should control the target weed(s) as effectively as the more resistance-prone partner.
- Herbicide use should be based on an integrated weed management program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical control methods), cultural (for example, higher crop seeding rates; precision fertilizer application method and timing to favour the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Monitor weed populations after herbicide application for signs of resistance development (for example, only one weed species on the herbicide label not controlled). If resistance is suspected, prevent weed seed production in the affected area if possible by an alternative herbicide from a different group. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.
- Have suspected resistant weed seeds tested by a qualified laboratory to confirm resistance and identify alternative herbicide options.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact Corteva Agriscience Canada Company at 1-800-667-3852.

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

TMTrademark of Corteva Agriscience and its affiliated companies.

All other products listed are registered trademarks of their respective companies.

020724

Label Code: CN-29985-009-E

Replaces: CN-29985-008-E

Specimen Notes:

Updated address, tank mix general statement and seed potato clarification



Tandem™ B

Herbicide

| | | |
|-------|---|-----------|
| GROUP | 4 | HERBICIDE |
|-------|---|-----------|

FOR SALE FOR USE ONLY IN THE PRAIRIE PROVINCES AND INTERIOR OF BRITISH COLUMBIA (INCLUDING THE PEACE RIVER REGION)

Tandem B Herbicide is a selective herbicide for post emergent control of annual broadleaved weeds including cleavers and certain ALS-resistant broadleaved weeds such as kochia in spring wheat, durum wheat and winter wheat. Tandem B Herbicide will also suppress hemp-nettle, common chickweed, and wild buckwheat.

COMMERCIAL (AGRICULTURAL)

READ THE LABEL AND BOOKLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN

ACTIVE INGREDIENT: fluroxypyr, present as 1-methylheptyl ester 333 g /L

Emulsifiable concentrate

REGISTRATION NO. 29965 PEST CONTROL PRODUCTS ACT

DANGER  **POISON**

**EYE AND SKIN IRRITANT
POTENTIAL SKIN SENSITIZER**

NET CONTENTS: 1 L – bulk

Corteva Agriscience Canada Company
Suite 240, 115 Quarry Park Rd. SE
Calgary, AB
T2C 5G9
1-800-667-3852

™ Trademark of Corteva Agriscience and its affiliated companies.

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN

Causes eye irritation. DO NOT get in eyes. May irritate the skin. Avoid contact with skin. Potential skin sensitizer.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12-hours.

DO NOT use in residential areas. Residential areas are defined as any use site where bystanders, including children, could be exposed during or after application. This includes homes, schools, parks, playgrounds, playing fields, public buildings, or any other area where the general public, including children, could be exposed.

Apply only when the potential for drift to areas of human habitation and areas of human activity such as houses, cottages, schools, and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.

PROTECTIVE CLOTHING AND EQUIPMENT

Wear a hat, coveralls over a long-sleeved shirt and long pants, socks, rubber boots and chemical-resistant gloves during spraying or application. In addition, wear goggles or face shield during mixing, loading, repair, and clean-up or when handling the concentrate. Applicators using a closed cab are not required to wear chemical-resistant gloves.

OPERATOR USE PRECAUTIONS

Wear freshly laundered clothing and clean protective equipment daily. At completion of spraying or end of the day take a shower immediately. Wash thoroughly with soap and water before eating, drinking, smoking or using the toilet. Remove contaminated clothing at the end of the work session and store and wash separately from household laundry using detergents and hot water before reuse. If herbicide penetrates clothing, remove immediately; then wash thoroughly and put on clean clothing. Throw away clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Avoid contact with eyes, skin and clothing. Avoid breathing spray mist.

PHYSICAL OR CHEMICAL HAZARDS

COMBUSTIBLE. Do not use or store near heat or open flame.

FIRST AID

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give **any** liquid to the person. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

TOXICOLOGICAL INFORMATION

THIS PRODUCT CONTAINS A PETROLEUM DISTILLATE. Vomiting may cause aspiration pneumonia. The decision of whether to induce vomiting or not should be made by the attending physician. If lavage is performed, suggest endotracheal and/or oesophageal control. Danger from lung aspiration of petroleum-based solvents must be weighed against toxicity when considering emptying the stomach. No specific antidote. Employ supportive care. Treatment should be based on the judgment of the physician in response to reactions of the patient.

AGRICULTURAL CHEMICAL

Do not ship or store with food, feeds, drugs, or clothing.

ENVIRONMENTAL PRECAUTIONS

TOXIC to non-target terrestrial plants and aquatic organisms. Observe spray buffer zones specified under DIRECTIONS FOR USE.

To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body.

This product contains an active ingredient and aromatic petroleum distillates which are toxic to aquatic organisms.

STORAGE

Store in original containers in a secure, dry heated storage. If product is frozen, bring to room temperature and agitate before use. Do not allow contamination of seeds, plants, fertilizers, or other pesticides. To prevent contamination, store this product away from food or feed. If containers are damaged or spill occurs, use the product immediately or contain the spill with absorbent materials and dispose of waste.

DISPOSAL

Recyclable Containers

Do not reuse this container for any purpose. This is a recyclable container and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial/territorial requirements.

Returnable Containers

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

Refillable Containers

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial/territorial regulatory agency. Contact the manufacturer and the provincial/territorial regulatory agency in case of a spill, and for clean-up of spills.

GENERAL INFORMATION

Tandem B Herbicide is a selective postemergent herbicide for the control of annual broadleaved weeds in spring wheat, durum wheat and winter wheat, not underseeded with legumes. Tandem B Herbicide offers a novel mode of action for controlling hard-to-kill annual broadleaved weeds such as kochia (including ALS resistant biotypes) and cleavers. Tandem B Herbicide will also provide suppression of hempnettle, wild buckwheat and chickweed.

MODE OF ACTION

Tandem B Herbicide is a systemic auxin-type herbicide and moves within the plant for control of exposed and underground plant tissues. The product controls weeds by disrupting normal plant growth patterns. Symptoms of weeds include epinasty (twisting of the stems) and swollen nodes.

GENERAL USE PRECAUTIONS

Sensitive Plants

Do not apply Tandem B Herbicide directly to, or otherwise permit it to come in direct contact with susceptible crops or desirable plants including alfalfa, edible beans, canola, flowers and ornamentals, lentils, lettuce, peas, potatoes, radishes, soybeans, sugar beets, sunflowers, tomatoes, or tobacco.

Non-Target Sites

Do not apply where proximity of susceptible crops or other desirable plants is likely to result in exposure to spray or spray drift. See Environmental Precautions section of the label.

Crop Rotation

Fields previously treated with Tandem B Herbicide can be seeded the following year to barley, canola, corn, dry beans, flax, lentils, mustard, oats, peas, potatoes, soybeans, sunflower, or wheat or fields can be summerfallowed.

Tank Mixtures

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact Corteva Agriscience Canada Company at 1-800-667-3852 or www.corteva.ca for information before applying any tank mix that is not specifically recommended on this label.

Spray Equipment Precaution

Do not apply through any type of irrigation system.

To Reduce Spray Drift:

- Use nozzles delivering higher volumes and coarser droplets.
- Use low pressures (200 to 275 kPa).
- Use 100 L/ha of spray solution.
- Spray when the wind velocity is 15 km/hr or less.
- Spot treatments should only be applied with a calibrated boom to prevent over-application.

Sprayer clean-out

To avoid injury to desirable plants, thoroughly clean equipment used to apply this product before re-use or using it to apply other chemicals.

1. Immediately after spraying, completely drain the sprayer tank. Any contamination on the outside of the spraying equipment should be removed by washing with clean water.
2. First rinse:
 - Spray the inside of tank with clean water and fill the sprayer with at least one tenth of the spray tank volume.
 - Agitate and circulate for 15 minutes, and flush through booms and hoses.
 - Remove end caps or open ball valves on the ends of each boom section, and flush solution through the boom ends to ensure there is no spray solution trapped between the boom end and the nozzles.
 - Drain tank completely.

3. Second rinse:
 - Fill the tank with clean water.
 - Add All Clear Spray Tank Decontaminator as per manufacturer's recommendations while filling the tank with clean water.
 - Agitate and then flush the boom and hoses with the cleaning solution. Top up with water making sure the tank is completely full. Allow to stand for 15 minutes with agitation. Flush the solution out of the spray tank through the spray booms. Remove end caps or open ball valves on the ends of each boom section, and flush solution through the boom ends to ensure there is no spray solution trapped between the boom end and the nozzles.
 - If possible, let the solution stand in the sprayer tank and booms for an extended period of time, overnight if possible.
 - After flushing the boom and hoses, drain tank completely.
 - Remove nozzles and screens and clean separately with a cleaning agent or an ammonia solution (100 mL in 10 L water).
4. Third rinse:
 - Rinse the tank with clean water and flush through the boom and hoses using at least one tenth of the spray tank volume.
 - Remove end caps or open ball valves on the ends of each boom section, and flush solution through the boom ends to ensure there is no spray solution trapped between the boom end and the nozzles.
 - Drain tank completely.

DIRECTIONS FOR USE

READ THE ENTIRE LABEL BEFORE USE. FAILURE TO FOLLOW LABEL INSTRUCTIONS MAY RESULT IN ERRATIC WEED CONTROL OR CROP DAMAGE. DO NOT APPLY TO CROPS UNDERSEEDING WITH LEGUMES.

As this product is not registered for the control of pests in aquatic systems, **DO NOT** use to control aquatic pests.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

Field sprayer application

DO NOT apply during period of dead calm. Avoid application of this product when winds are gusty. DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) medium classification. Boom height must be 60 cm or less above the crop or ground.

Aerial application

DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT apply when wind speed is greater than 16 km/h at flying height at the site of application. DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) medium classification. Reduce drift caused by turbulent wingtip vortices. Nozzle distribution along the spray boom length MUST NOT exceed 65% of the wing or rotor span.

Crops Registered

spring wheat
durum wheat
winter wheat

TANDEM B HERBICIDE ALONE

Apply Tandem B Herbicide for postemergent control of cleavers and kochia, including ALS resistant kochia and other broadleaved weeds infesting spring wheat, durum wheat and winter wheat. See following table for weeds controlled/suppressed and applications rates.

Weeds Controlled or Suppressed By Tandem B Herbicide Alone

| Weeds Controlled | Tandem B Herbicide Rate (L/ha) |
|--------------------------------|--------------------------------|
| chickweed* (up to 8 cm) | 0.41 |
| cleavers (1-6 whorls) | 0.21 |
| cleavers (1-8 whorls) | 0.31 |
| kochia* (2-8 leaf) | 0.31 |
| round-leaved mallow (1-6 leaf) | 0.41 |
| volunteer flax (1-12 cm) | 0.31 |
| Weeds Suppressed | Tandem B Herbicide Rate (L/ha) |
| hempnettle (2-6 leaf) | 0.41 |
| stork's-bill (1-8 leaf) | 0.31 |
| wild buckwheat (1-4 leaf) | 0.21 |

*Including biotypes resistant to Group 2 herbicides that inhibit the ALS enzyme.

NOTE: Tandem B Herbicide activity is influenced by weather conditions. Optimum activity requires active crop and weed growth. The temperature range for optimum activity is 12°C to 24°C. Reduced activity will occur when temperatures are below 8°C or above 27°C. Frost before application (3 days) or shortly after (3 days) may reduce weed control and crop tolerance. Weed control may be reduced during stress conditions, e.g. drought, heat, or cold stress, or if weeds have initiated flowering, or if heavy infestations exist.

APPLICATION DIRECTIONS**Application Timing**

Apply to actively growing spring wheat and durum wheat from the 2-leaf crop growth stage up to and including initiation of stem elongation. Apply to winter wheat in the spring from the 3-tiller stage to just before the flag leaf stage. Apply when weeds are actively growing. Best results are obtained from applications made to seedling weeds. Only weeds emerged at the time of treatment will be controlled. Extreme growing conditions such as drought or near freezing temperature prior to, at and following time of application may reduce weed control and increase the risk of crop injury at all stages of growth. Foliage that is wet at the time of application may decrease control. Tandem B Herbicide applications are rainfast within 1 hour after application. Under conditions of low crop and high weed density, control may be reduced.

Preharvest/Grazing Intervals (Tandem B Herbicide applied alone)

- Livestock may be grazed on treated crops 3 days following application.
- Do not harvest the treated crop within 60 days after application.

APPLICATION DIRECTIONS (GROUND AND AERIAL APPLICATIONS)**(1) Ground Application**

Using ground equipment, apply Tandem B alone or with tank mix partners as a broadcast treatment. Apply Tandem B alone at the recommended rates in a spray volume of 50-100 L water/ha.

(2) Aerial Application**Do not use human flaggers.**

Use Tandem B alone or tank mixed with a product registered for aerial application as a broadcast treatment. Apply the recommended rates of Tandem B and tank mix partner as indicated elsewhere on this label. Apply in a spray volume of 30-50 L water/ha.

Apply only by fixed-wing or rotary aircraft equipment which has been functionally and operationally calibrated for the atmospheric conditions of the area and the application rates and conditions of this label.

Label rates, conditions and precautions are product specific. Read and understand the entire label before opening this product. Apply only at the rate recommended for aerial application on this label. Where no rate for aerial application appears for the specific use, this product cannot be applied by any type of aerial equipment.

Ensure uniform application. To avoid streaked, uneven or overlapped application, use appropriate swath marking devices.

Use Precautions

Apply only when meteorological conditions at the treatment site allow for complete and even coverage. Apply only under conditions of good practice specific to aerial application as outlined in the National Aerial Pesticide Application Manual, developed by the Federal/Provincial/Territorial Committee on Pest Management and Pesticides.

Operator Precautions

Do not allow the pilot to mix chemicals to be loaded onto the aircraft. Loading of premixed chemicals with a closed system is permitted.

It is desirable that the pilot have communication capabilities at each treatment site at the time of application.

The field crew and the mixer/loaders must wear chemical resistant gloves, coveralls and goggles or face shield during mixing/loading, cleanup, and repair. Follow the more stringent label precautions in cases where the operator precautions exceed the generic label recommendations on the existing ground boom label.

All personnel on the job site must wash hands and face thoroughly before eating and drinking. Protective clothing, aircraft cockpit and vehicle cabs must be decontaminated regularly.

Product Specific Precautions

Read and understand the entire label before opening this product. If you have questions, call the manufacturer at 1-800-667-3852 or obtain technical advice from the distributor or your provincial agricultural representative. Application of this specific product must meet and/or conform to the following:

- The interaction of many equipment-and-weather-related factors determine the potential for spray drift. Users are responsible for considering all these factors when making decisions.

The following drift management requirement must be followed to avoid off-target drift movement from aerial applications: Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.

TANK-MIX COMBINATION – TANDEM A HERBICIDE + TANDEM B HERBICIDE OR TANDEM GODRI A HERBICIDE + TANDEM B HERBICIDE

For control of a wide spectrum of grass and broadleaf weeds in spring and durum wheat, apply Tandem A tank mixed with Tandem B or Tandem GoDRI A tank mixed with Tandem B. Apply post-emergence when weeds are actively growing: grass weeds up to the 4 leaf, 2 tiller stage and broadleaf weeds as indicated in the following table. Best results are obtained from applications made to seedling weeds. Apply to actively growing spring or durum wheat from the 3-leaf stage to the first node stage. Apply to winter wheat in the spring in the 2-7 leaf, 4 tiller stage. Extreme growing conditions such as drought or near freezing temperature prior to, at, or following time of application may reduce weed control and increase the risk of crop injury at all stages of growth. Under conditions of low crop and high weed density, control may be reduced.

Preharvest/Grazing Intervals (Tandem A and Tandem B or Tandem GoDRI A and Tandem B)

- Livestock may be grazed on treated crops 7 days following application.
- Do not harvest the treated crop within 60 days after application.

Rates of Application

I. Low Rate: To control the weeds listed below, apply a uniform spray containing Tandem A at 375 mL/ha tank mixed with Tandem B at 0.21 L/ha or Tandem GoDRI A at 52 g/ha tank mixed with Tandem B at 0.21 L/ha.

Weeds Controlled

| | |
|--------------------------------------|-----------------------|
| wild oats (up to 4 leaf, 2 tiller) * | cleavers (1-6 whorls) |
|--------------------------------------|-----------------------|

*Under low wild oat populations (<75 plants/m²)

Refer to the tank-mix combinations table for a complete list of additional weeds controlled when tank mixed with 2,4-D or MCPA.

II. High Rate: To control the weeds listed below, apply a uniform spray containing Tandem A at 500 mL/ha tank mixed with Tandem B at 0.31 L/ha or Tandem GoDRI A at 70 g/ha tank mixed with Tandem B at 0.31 L/ha.

Weeds Controlled

| | |
|--|--|
| barnyard grass (1-5 leaf) | kochia* (2-8 leaf) |
| chickweed, common (up to 10 cm) | red root pigweed (1-8 leaf) |
| cleavers (1-8 whorls) | round leaved mallow (up to 6 leaf stage, <10 cm in size) |
| corn spurry (up to 2 whorl stage, <10cm in height) | shepherd's purse (up to 30 cm tall) |
| cow cockle (up to 8 leaf) | smartweed (lady's thumb, 1-5 leaf) |
| flixweed (up to 10 cm) | stinkweed (up to 30 cm tall) |
| foxtail, yellow (1-5 leaf) | volunteer canola** (1-6 leaf) |
| hemp nettle (1-8 leaf) | volunteer flax (1-12 cm) |
| Japanese brome (1-6 leaf) | wild oats (up to 4 leaf, 2 tiller) |

*Including biotypes resistant to Group 2 herbicides that inhibit the ALS enzyme.

** Will not control volunteer Clearfield® canola.

Weeds Suppressed

| | |
|--|------------------------------------|
| brome, downy (2-6 leaf, 4 tillers) | Russian thistle (up to 10 cm tall) |
| Canada thistle (up to 30 cm tall, prebud) | storks-bill (1-8 leaf) |
| dandelion (seedlings and over-wintered rosettes less than or equal to 20 cm) | wild buckwheat (1-4 leaf) |
| foxtail, green (1-5 leaf) | |

TANK-MIX COMBINATIONS – TANDEM A + TANDEM B + OTHER HERBICIDES OR TANDEM GODRI A + TANDEM B + OTHER HERBICIDES

When tank mixing Tandem A + Tandem B or Tandem GoDRI A + Tandem B the addition of other herbicides will provide control of additional weeds. For products which may be tank mixed with Tandem A + Tandem B or Tandem GoDRI A + Tandem B and the additional weeds controlled, see following table.

Tank-Mix Combinations with Tandem A (375 mL/ha) + Tandem B (0.21 L/ha) or Tandem A (500 mL/ha) + Tandem B (0.31 L/ha) or Tandem GoDRI A (52 g/ha) + Tandem B (0.21 L/ha) or Tandem GoDRI A (70 g/ha) + Tandem B (0.31 L/ha)

| Herbicide Tank-Mix Partner | Crops Registered | Rate of Tank-Mix Partner | Additional Weeds Controlled |
|-----------------------------------|--|---------------------------------|--|
| MCPA Ester* | spring wheat durum wheat winter wheat | Up to 560 g ae/ha | burdock (before the 4-leaf stage), cocklebur, lamb's-quarters, mustards (except dog and tansy), prickly lettuce, plantain, ragweeds, Russian pigweed, vetch, wild buckwheat, wild radish, wild (annual) sunflower. *Consult MCPA Ester labels for weed stages and rates |
| Curtail™ M♦ (ground only) | spring wheat durum wheat | 1.5 L/ha | Canada thistle ¹ (low infestations), burdock, cocklebur, field horsetail ² , lamb's quarters, wild mustard, plantain ² , prickly lettuce, wild radish, ragweed, annual sowthistle (1-5 leaf) ³ , annual sunflower, volunteer sunflower, vetch ♦Consult Curtail M labels for weed stages and rates 1 Season long control, with some re-growth in fall 2 Top growth only 3 Only controlled with the high rate of Tandem A+ Tandem B Tandem or GoDRI A + Tandem B |
| 2,4-D Ester* | spring wheat, durum wheat winter wheat | Up to 420 g ae/ha | Susceptible weeds such as: bluebur, burdock, cocklebur, goat's-beard, lamb's-quarters, mustard (except dog and green tansy), plantain, prickly lettuce, ragweeds, sunflower (annual), sweet clover, volunteer canola, wild buckwheat, wild radish *Consult 2,4-D Ester labels for weed stages and rates |

SPRAY BUFFER ZONES

A spray buffer zone is NOT required for:

- uses with hand-held application equipment permitted on this label.

The spray buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands) and sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs, and wetlands) and estuarine/marine habitats.

| Method of application | Crop | | Spray Buffer Zones (meters) Required for the Protection of: | | | | |
|-----------------------|---|-------------|---|------------------|-------------------------------------|------------------|---------------------|
| | | | Freshwater Habitat of Depths: | | Estuarine/marine Habitat of Depths: | | Terrestrial habitat |
| | | | Less than 1 m | Greater than 1 m | Less than 1 m | Greater than 1 m | |
| Field sprayer | Spring wheat, durum wheat, winter wheat | | 1 | 0 | 1 | 1 | 3 |
| Aerial | Spring wheat, durum wheat, winter wheat | Fixed Wing | 5 | 0 | 1 | 1 | 95 |
| | | Rotary Wing | 3 | 0 | 1 | 1 | 80 |

When tank mixes are permitted, consult the labels of the tank-mix partners, and observe the largest (most restrictive) spray buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.

The spray buffer zones for this product can be modified based on weather conditions and spray equipment configuration by accessing the Spray Buffer Zone Calculator on the Pesticides portion of the Canada.ca website.

RESISTANCE MANAGEMENT RECOMMENDATIONS

For resistance management, please note that Tandem B Herbicide is a Group 4 herbicide. Any weed population may contain or develop plants naturally resistant to Tandem B Herbicide and other Group 4 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

- Where possible, rotate the use of Tandem B Herbicide or other Group 4 herbicide within a growing season (sequence) or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group when such use is permitted. To delay resistance, the less resistance-prone partner should control the target weed(s) as effectively as the more resistance-prone partner.
- Herbicide use should be based on an integrated weed management program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical control methods), cultural (for example, higher crop seeding rates; precision fertilizer application method and timing to favour the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Monitor weed populations after herbicide application for signs of resistance development (for example, only one weed species on the herbicide label not controlled). If resistance is suspected, prevent weed seed production in the affected area, if possible, by an alternative herbicide from a different group. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields and planting clean seed.
- Have suspected resistant weed seeds tested by a qualified laboratory to confirm resistance and identify alternative herbicide options.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact Corteva Agriscience Canada Company at 1-800-667-3852 or at www.corteva.ca.

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

™ Trademark of Corteva Agriscience and its affiliated companies.

040424

Label Code: CN-29965-009-E

Replaces: CN-29965-008-E

Specimen Notes:

Updated tank mix general statement and address




Water Conditioner

BEFORE HANDLING OR USING THIS PRODUCT, SEE YOUR EMPLOYER AND READ THE CURRENT MATERIAL SAFETY DATA SHEET

ACTIVE INGREDIENT

Ammonium Hydroxide: 9.75%

DANGER



CAUSES SEVERE EYE BURNS•CAUSES SKIN BURNS•HARMFUL IF INHALED; MAY CAUSE LUNG INJURY•MAY CAUSE RESPIRATORY TRACT IRRITATION•ASPIRATION HAZARD. CAN ENTER LUNGS AND CAUSE DAMAGE.

Keep out of reach of children. Do not get in eyes, on skin, on clothing. Do not swallow. Avoid breathing vapor or mist. Wash thoroughly after handling. Keep container closed. Use with adequate ventilation.

FIRST AID

IN CASE OF EYE CONTACT, immediately flush eyes with plenty of water for at least 30 minutes. Remove contact lenses, if worn. Get medical attention immediately.

IN CASE OF SKIN CONTACT, immediately flush skin with plenty of water for at least 15 minutes while removing contaminated clothing and shoes. Get medical attention immediately. Wash clothing before reuse. Destroy contaminated shoes.

IF SWALLOWED, call a physician and/or transport to a medical facility immediately. Do NOT induce vomiting unless directed to do so by medical personnel. Never give anything by mouth to an unconscious person.

IF INHALED, remove to fresh air. If not breathing, give artificial respiration. If breathing is difficult, oxygen should be administered by qualified personnel. Get medical attention.

TOXICOLOGICAL INFORMATION

No specific antidote. Employ supportive care. Treatment should be based on judgement of the physician in response to reactions of the patient.

STORAGE

Keep in original container during storage. Store product in cool, dry, well ventilated place away from seed, feed, fertilizer, or other pesticides. Do not store in aluminum.

DISPOSAL

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

MIXING INSTRUCTIONS

1. Fill sprayer tank 1/2 full of water and continue filling while adding products to the tank.
2. Start sprayer tank agitation and continue agitating throughout filling process.
3. Add 1.5 L of Water Conditioner per jug of Simplicity to be mixed in the spray tank.
4. Follow the mixing instructions on the product label.

Net Contents: 0.40 gal/1.5 L

Corteva Agriscience Canada Company

Suite 240, 115 Quarry Park Rd. SE
Calgary, AB
T2C 5G9
1-800-667-3852

™Trademark of Corteva Agriscience and its affiliated companies.

For product information, call 1-800-667-3852 or visit our web site at www.corteva.ca.

181223

Label Code: CN-Water Cond-004-E

Replaces: CN-Water Cond-003-E

Label Notes:

Update address