THIS RECOMMENDATION IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE EPA.



EPA REG. NO. 84876-2

2(ee) Recommendation

FOR DISTRIBUTION AND USE ONLY IN California

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING

This Product Bulletin Contains New or Additional Directions For Use Which Are Recommended By Sym-Agro, Inc.Under FIFRA Section 2(ee)

CROP	PEST	RATE OF APPLICATION
Almonds	Hull Rot (<i>Monilinia</i> spp.)	20.4– 45.7 fl oz/A (0.34-0.75 lb ai/A)

DIRECTIONS FOR USE:

Monilinia spp.: For control of Hull Rot apply ProBlad® Verde prior to onset of disease development. Monilinia Hull Rot is best managed with treatments 3-4 weeks prior to hull split. A second application should be made 7-14 days after initial hull split application to assure protection through hull split growth stage.

General: Increase spray volume as growth increases in order to provide thorough coverage of foliage and fruit for optimum disease control.

RESTRICTIONS:

Do not make more than two sequential applications of ProBlad® Verde Fungicide before alternating to a labeled fungicide with a different mode of action.

Do not make more than 5 applications of ProBlad® Verde Fungicide per harvest cycle.

PHI: 1 Day

ALL APPLICABLE DIRECTIONS, RESTRICTIONS, WORKER PROTECTION STANDARDS REQUIREMENTS AND PRECAUTIONS ON THE EPA REGISTERED PROBLAD VERDE® FUNGICIDE (EPA REG. NO. 84876-2) LABEL MUST BE FOLLOWED. THIS RECCOMENDATION MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.



Sym-Agro,Inc. PO Box 71 Visalia, CA 93291 ProBlad is a trademark of CEV S.A.

ProBlad Verde 2ee Almond, Hull Rot 04-06-2020