

Bayer CropScience LP P.O. Box 12014 2 T.W. Alexander Drive Research Triangle Park, North Carolina 27709 1-866-99BAYER (1-866-992-2937)

Leverage® 360 Insecticide

For the Suppression of Soybean Stem Borer (Dectes Stem Borer) Adults on Soybean

EPA Reg. No. 264-1104

FIFRA 2(ee) Recommendation

RESTRICTED USE PESTICIDE

Due to Toxicity to Fish and Aquatic Organisms

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

FOR DISTRIBUTION AND USE ONLY IN THE STATE OF NEBRASKA

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its label.

SOYBEAN		
PEST SUPPRESSED	Product Rate fluid ounces/Acre	Method
Adults of Soybean Stem Borer (Dectes Stem Borer)	2.8	Apply as a foliar spray, using sufficient carrier volume for complete coverage of stems and foliage. Applications should only target the adult life stage upon overwintering emergence from stems; applications made after this time or targeting larval or pupal life stages within the soybean stem will not provide any activity.

RESTRICTIONS

Pre-Harvest Interval (PHI) for seed: **21 days**. Dry vines (hay) and green forage may be fed **15 days** after last application Minimum interval between applications: **7 days**

Maximum LEVERAGE 360 Insecticide allowed per year: **9.0 fluid ounces/Acre** (0.07 lb Al/A β -cyfluthrin, 0.14 lb Al/A imidacloprid)

Maximum β-cyfluthrin allowed in all forms per crop season: 0.088 lb Al/A Maximum cyfluthrin allowed in all forms per crop season: 0.175 lb Al/A

Maximum β-cyfluthrin plus cyfluthrin allowed in all forms per crop season: 0.175 lb Al/A

ADDITIONAL USE DIRECTIONS

LEVERAGE 360 Insecticide is to be applied through properly calibrated ground, aerial or chemigation application equipment.

For <u>MEDICAL</u> And <u>TRANSPORTATION</u> Emergencies <u>ONLY</u> Call 24 Hours A Day 1-800-334-7577. For <u>PRODUCT USE</u> Information Call 1-866-99BAYER (1-866-992-2937).

As with any crop-protection product, always read and follow label directions. For additional information call toll-free 1-866-99BAYER (1-866-992-2937).