



Section 24(c) Special Local Need Label

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF DELEWARE

**Dual Magnum® Herbicide
For Weed Control in Lima Bean**

**EPA Reg. No. 100-816
EPA SLN No. DE-21-0001**

**This label expires and must not be distributed or used in accordance with this SLN
registration after December 31, 2026**

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This label must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA-registered label.
- The risk of crop injury increases when adjuvants (e.g., non-ionic surfactants, crop oils, etc.), nitrogen sources (e.g., AMS, UAN), fertilizers or other pesticides are applied with Dual Magnum.

Postemergence Applications

If a preplant, preplant incorporated or preemergence application of Dual Magnum was made to the crop, a postemergence application may be made provided the total amount of Dual Magnum applied does not exceed 2.0 pt/A. Refer to the Dual Magnum label for preplant, preplant incorporated and preemergence application use directions.

Apply Dual Magnum postemergence only after the first trifoliate stage of plant growth. Application to plants with less than one trifoliate can result in unacceptable crop injury. When applied broadcast over the top, crop injury in the form of leaf spotting and speckling may be observed, especially with rates greater than 0.75 pt/A

Precautions:

- 1) Application to wet plants or when conditions are extremely hot or humid may result in increased risk of crop injury.
- 2) The risk of crop injury is greater on lighter textured soils and with higher use rates, especially when coupled with heavy rains or when excessive multiple irrigations occur within 5 days of application.
- 3) Dual Magnum will not control emerged weeds; thus, other control tactics that provide weed control prior to Dual Magnum being activated are required.

- 4) All cultivars have not been tested for tolerance. Experiment on a limited basis until on-farm confidence in these use patterns and rates are obtained.
- 5) Dual Magnum should be applied only with water as the carrier or unacceptable crop injury may occur.

Restrictions:

- 1) Do not apply more than 2.0 pt/A of Dual Magnum total per season.
 - 2) Do not apply from the time of soil cracking until the first trifoliate is fully expanded.
 - 3) Do not apply this product through an irrigation system other than a center-pivot irrigation system.
 - 4) Do not harvest crops within 50 days of a Dual Magnum application.
 - 5) Do not feed animals with plant material treated with a post-emergence application of Dual Magnum.
-

Dual Magnum® and the Syngenta logo are trademarks of a Syngenta Group Company.

©2021 Syngenta

24(c) Registrant:
Syngenta Crop Protection, LLC
P. O. Box 18300
Greensboro, NC 27419-8300

Label Code: DE0816031AA0421