

FIFRA SECTION 2(ee) RECOMMENDATION FOR USE IN CALIFORNIA


FUNGI-PHITE®
EPA REG. NO.: 73771-5
FOR USE IN CALIFORNIA**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product inconsistent with its labeling. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label. This recommendation must be in the possession of the user at the time of pesticide application.

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA.

POME FRUIT CROPS, such as:

Apple, Crabapple, Loquat, Mayhaw, Pear, Asian Pear and Quince

DISEASE**	APPLICATION METHOD	APPLICATION RATE	APPLICATION PROGRAM
†Suppression of: <i>Botryosphaeria</i> spp. (including <i>dothidea</i> ,) <i>Pseudomonas</i> spp. (including <i>syringae</i>), <i>Xanthomonas</i> spp. (including <i>campestris</i>), and <i>Eutypa lata</i>	Foliar Spray		
	Aerial:	Apply 1-3 quarts/acre (2-7 L/ha) in a minimum of 20 gallons/acre (190 L/ha) of water.	Begin application after plants are established and from first lead flush when conditions favor disease development. Disease Prevention/Low Pressure Program* : Apply lower rate at 2-4 week intervals. Do not apply more than 4 times per crop cycle. Disease Control/High Pressure Program* : Apply higher rate at 1-3 week intervals. Do not apply more than 4 times per crop cycle.
	Ground:	Apply 1-3 quarts/acre (2-7 L/ha) in a minimum of 100 gallons/acre (950 L/ha) of water.	
	Chemigation Overhead:	Apply 2-3 quarts/acre (5-7 L/ha) in a minimum of 1,000 gallons/acre (9,350 L/ha) of water.	Apply with normal irrigation schedule. Follow Disease Programs* as stated above. Do not apply more than 6 times per crop cycle. Low Disease Pressure* : Apply lower rate at the first onset of the disease. Repeat applications at 1-2 week intervals. Do not apply more than 6 times per crop cycle.
	Foliar Spray	Apply 1-2.5 quarts/acre (3-7 L/ha) in a minimum of 100 gallons/acre (950 L/ha) of water.	High Disease Pressure* : Apply higher rate at the first onset of the disease. Repeat applications at 7-10 days intervals. Do not apply more than 6 times per crop cycle. Apply during active plant growth.
	Trunk Spray	Mix 1-2 quarts with (1-2 L) in 5 gallons (20 L) of water.	Use higher rates when lesions are present. Clean the wound site and apply on and around the lesions using enough spray volume to thoroughly wet the lesions. In the absence of lesions, use the lower rate and apply to the trunk from the soil line to 2 feet up the trunk. Apply one time in the Spring, Summer, and Fall.

* Check with your local Extension Agent or Crop Consultant if you are unsure about disease prevention, control or severity/pressure.

** See the label booklet for Fungi-Phite for a full list of diseases pertaining to this crop group.

† Fungi-Phite has suppressive properties on diseases caused by certain bacteria or fungi. When applying Fungi-Phite for disease suppression, use it in combination with another registered bactericide or fungicide registered for the same crops listed on the label.

Read the "Conditions of Sale and Limitations of Warranty and Liability" in the label booklet for Fungi-Phite before using.