

FIFRA Section 2(ee) Recommendation



FOR DISTRIBUTION AND USE ONLY IN THE STATE OF TEXAS

BELAY[®] **INSECTICIDE**

EPA Reg. No. 59639-150

BELAY[®] INSECTICIDE FOR THE CONTROL OF REDBANDED STINK BUG (*Piezodorus guildinii* Westwood) IN SOYBEANS

This FIFRA Section 2(ee) recommendation expires December 31, 2025.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This 2(ee) recommendation contains directions for use which do not appear on the package label and must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions and precautions on the EPA registered label when using *Belay* Insecticide alone or in tank mix combinations. The most restrictive labeling applies when using a tank mix.

BELAY INSECTICIDE APPLICATION TO SOYBEANS

	FOLIAR APPLICATION		
Pest	<i>Belay</i> Insecticide Application Rate	Mixture partner application rate	Instructions
Redbanded stink bug	2.0 fl oz/A (0.03 lb ai/A)	Bifenthrin 0.06 to 0.10 lbai/A	For the control of adults and immature stages of redbanded stinkbug. Bifenthrin will aid by delivering a faster knockdown of adults.
	2.0 fl oz/A (0.03 lb ai/A)	Acephate 0.75 to 1.0 lbai/A	For the control of adults and immature stages of redbanded stinkbug. Acephate will aid by delivering a faster knockdown of adults.

	4.0- 6.0 fl oz/A		Apply Belay for the control of redbanded stink bugs when adults are first observed in the field.
	RESTRICTIONS <ul style="list-style-type: none"> • Do not make foliar applications of Belay insecticide in fields treated with a neonicotinoid insecticide seed treatment(s) within 45 days after planting. • Regardless of formulation or type of application method, do not apply more than 0.2 lb active ingredient clothianidin per acre per year. • Do not apply more than 0.3 lb active ingredient bifenthrin per acre per year. • Do not apply more than 1.5 lb active ingredient acephate per acre per year • Do not apply treatments less than 7 days apart for the application of Belay alone or the acephate mixture. • Do not apply treatments less than 30 days apart for the bifenthrin mixture. • Do not apply within 21 days of harvest. • <i>Belay</i> Insecticide must not be applied during bloom or when bees are foraging. • Do not feed or allow livestock to graze on cover crops from treated orchards. 		

Please contact Valent U.S.A. LLC at 1-800-6-VALENT (682-5368) to determine if this product is registered in your state.

© 2020 by Valent U.S.A. LLC

Registrant: Valent U.S.A. LLC
P.O. Box 8025
Walnut Creek, CA 94596-8025

Belay is a registered trademark of Sumitomo Chemical Company, Ltd.