

TECHNICAL INFORMATION BULLETIN
FOR THE USE OF GRANDEVO® WDG ON SWEET POTATO FOR MANAGEMENT OF NEMATODES

FIFRA Section 2(ee) Recommendation

This recommendation is made as permitted by Section 2(ee) of FIFRA, as amended, and has not been submitted to or approved by the U. S. Environmental Protection Agency.

FOR USE IN ALL STATES WHERE CROP AND PEST(S) EXIST AND WHERE 2(EE) RECOMMENDATIONS ARE
RECOGNIZED.

GRANDEVO® WDG
EPA Reg. No. 84059-27

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Crop	Target Nematodes	Application Method	Product Use Rate per Application	Application Instructions
Sweet Potato	Root knot nematode (<i>Meloidogyne incognita</i>); stubby root nematode (<i>Paratrichodorus</i> sp.); ring nematode (<i>Criconeoides xenoplax</i>)	Soil through drip irrigation	3 lb/Acre	Minimum of two applications, the first application at two weeks after transplanting, and the second application at four to six weeks after transplanting. The in-season use of Grandevo WDG does not eliminate the need for a pre-plant fumigation for nematode control
GRANDEVO® WDG has a pre-harvest interval (PHI) of 0 days. Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 hours.				

All applicable directions, restrictions and precautions on the EPA-registered label are to be followed.

This FIFRA Section 2(ee) recommendation contains new or additional directions for use of this product, which may not appear on the package label. Read and carefully observe the cautionary statements plus all the other information appearing on the product label.

This recommendation must be in the possession of the user at the time of pesticide application.

GRANDEVO® WDG is a registered trademark of Marrone Bio Innovations, Inc.
© Copyright Marrone Bio Innovations, Inc., 2020
Marrone Bio Innovations, Inc.
1540 Drew Ave.
Davis, California 95618
Telephone (530) 750-2800
www.marronebioinnovations.com
20200214A