

Product Bulletin

Corteva Agriscience LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Intrepid Edge®

EPA Reg. No. 62719-666

2(ee) Recommendation†

For Distribution and Use Only in the State of Texas

This product bulletin expires December 31, 2027 and must not be used or distributed after this date.

For Control of Walnut Husk Fly in Walnut

ATTENTION

† This recommendation is made as permitted under FIFRA 2(ee) and has not been submitted to or approved by the EPA.

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- Read and follow all applicable directions for use, precautions, and limitations on the product label attached to the container for this product.

Directions for Use

Refer to the Intrepid Edge® product label for General Use Precautions, Mixing Directions, Application Directions, and Restrictions.

Application Rate: The amount of Intrepid Edge applied per acre will depend upon tree size and pest infestation level. Use a lower rate in the rate range for light infestations and/or smaller trees and a higher rate in the rate range for heavy infestations and/or larger trees.

Ground Application: Apply in a minimum of 50 gpa by conventional ground equipment. For best results, use 100 to 200 gpa. Use a spray volume that assures uniform coverage of the infested crop.

Aerial Application: Apply in a minimum of 10 gpa. This method may result in reduced efficacy if the size of the tree or density of the foliage prohibits thorough, uniform coverage of the entire tree canopy.

Resistance Management: Do not make more than two consecutive applications of Group 5 or 18 insecticides per crop. If additional treatments are required after two consecutive applications of Group 5 or 18 insecticides, rotate to another class of effective insecticide for at least one application. Consult your Dow AgroSciences representative, extension specialist, certified crop advisor, or state agricultural experiment station for information on alternative effective products to use in your area. Avoid applying Intrepid Edge and other Group 5 and 18 insecticides to consecutive generations of all target insect pests.

Pest	Intrepid Edge® (fl oz/acre)
Walnut husk fly	12 – 14

Restrictions:

- **Preharvest Interval:** Do not apply within 7 days of harvest.
- Do not apply more than 18 fl oz of Intrepid Edge per acre per application.
- Do not apply more than a total of 51 fl oz of Intrepid Edge® (1 lb ai methoxyfenozide, 0.2 lb ai spinetoram) per acre per season.
- **Minimum Treatment Interval:** Do not make applications less than 14 days apart.

- **Maximum Number of Applications:** Do not make more than 4 applications per year.
- Do not make more than two consecutive applications.

®™Trademarks of Corteva Agriscience and its affiliated companies

R911-020

Approved: 02/28/2024

Replaces R911-006