

FIFRA Section 2(ee) Recommendation



FOR DISTRIBUTION AND USE ONLY IN THE STATES OF ALABAMA, ARKANSAS, FLORIDA, GEORGIA, LOUISIANA, MISSISSIPPI, NORTH CAROLINA, OKLAHOMA, SOUTH CAROLINA, TENNESSEE, VIRGINIA



EPA Reg. No. 59639-99



EPA Reg. No. 59639-221

FOR USE IN TANK MIXES TO CONTROL ANNUAL BROADLEAF AND GRASS WEEDS IN COTTON

This FIFRA Section 2(ee) Recommendation Expires December 31, 2028.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This recommendation is made as permitted by Section 2(ee) of FIFRA, as amended, and has not been submitted to or approved by the U.S. Environmental Protection Agency. This 2(ee) recommendation contains directions for use which do not appear on the package label and must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label when using Valor[®] SX Herbicide or Valor[®] EZ Herbicide². Always read and follow all label directions when using any pesticide alone or in tank mix combinations. The most restrictive labeling applied when using a tank mix.

APPLICATION TO COTTON

HERBICIDE TANK MIX	APPLICATION RATES	APPLICATION INSTRUCTIONS— POST DIRECTED AND LAYBY USE DIRECTIONS
<i>Valor</i> SX Herbicide or <i>Valor</i> EZ Herbicide ² + Glufosinate ammonium or Diuron or Paraquat or MSMA + Diuron	2.0 oz/A or 2.0 fl oz/A + LUR ¹ or LUR ¹ or LUR ¹ or LUR ¹ + LUR ¹	<ul style="list-style-type: none">For postemergence weed control apply <i>Valor</i> SX Herbicide or <i>Valor</i> EZ Herbicide² through a hooded or shielded sprayer or at layby in combination with glufosinate ammonium, diuron, paraquat, or MSMA + diuron to assist in the control of annual broadleaf and grass weeds.To ensure thorough coverage in hooded, shielded, and layby applications, use a minimum of 15 gallons spray solution per treated acre. Use a minimum of 20 gallons per treated acre under heavy weed pressure.Add an agronomically approved non-ionic surfactant. Do not use crop oil concentrates, methylated seed oils, organo-silicone surfactants or products containing these ingredients.Hooded and shielded applications may be applied after cotton has reached a minimum of 6 inches in height.Layby applications may be made once cotton has reached a minimum of 16 inches in height. Applications must be directed to the lower 2 inches of the cotton stem to avoid injury.<i>Valor</i> SX Herbicide or <i>Valor</i> EZ Herbicide² tank mixtures with glufosinate must be made in Liberty Link[®] cotton only.
<ul style="list-style-type: none">Do not apply more than 2 oz of <i>Valor</i> SX Herbicide or 2 fl oz of <i>Valor</i> EZ Herbicide² per acre per application.Do not make more than 2 applications of <i>Valor</i> SX Herbicide or <i>Valor</i> EZ Herbicide² per acre per year.Do not make a sequential application of <i>Valor</i> SX Herbicide or <i>Valor</i> EZ Herbicide² within 30 days of the first application.Do not apply more than 4 oz of <i>Valor</i> SX Herbicide or 4 fl oz of <i>Valor</i> EZ Herbicide² per acre per year.Do not apply within 60 days of harvest.		

¹ Labeled Use Rate.

Please contact Valent U.S.A. LLC at 800-6-VALENT (682-5368) to determine if this product is registered in your state.

©2024 Valent U.S.A. LLC

Registrant: Valent U.S.A. LLC
P.O. Box 5075
San Ramon, CA 94583

Valor is a registered trademark of Valent U.S.A. LLC and Liberty Link is a registered trademark of Bayer.