

FIFRA SECTION 2(ee) RECOMMENDATION FOR USE IN CALIFORNIA



FUNGI-PHITE®
EPA REG. NO.: 73771-5

FOR USE IN CALIFORNIA

DIRECTIONS FOR USE

It is a violation of Federal law to use this product inconsistent with its labeling. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label. This recommendation must be in the possession of the user at the time of pesticide application.

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA.

KIWI, OLIVES AND TREE NUT CROPS, such as:

Almond, Beech Nut, Brazil Nut, Butternut, Cashew, Chestnut, Chinquapin, Filbert, Hazelnuts, Hickory Nut, Macadamia Nut, Pecan, Pistachio, and Walnut

DISEASE**	APPLICATION METHOD	APPLICATION RATE	APPLICATION PROGRAM
† Suppression of: <i>Botryosphaeria</i> spp. (including <i>dothidea</i>), <i>Pseudomonas</i> spp. (including <i>syringae</i>), <i>Xanthomonas</i> spp. (including <i>campestris</i>), and <i>Eutypa lata</i>	Foliar Spray		
	Aerial:	Apply 1-2 quarts/acre (2-5 L/ha) in a minimum of 20 gallons/acre (190 L/ha) of water.	Begin application during active plant growth after plants are established and conditions favor disease development. Disease Prevention/Low Pressure Program* : Apply lower rate at 3 monthly intervals. Do not apply more than 4 times per crop cycle. Disease Control/High Pressure Program* : Apply higher rate at monthly intervals. Do not apply more than 4 times per crop cycle.
	Ground:	Apply 1-2 quarts/acre (2-5 L/ha) in a minimum of 100 gallons/acre (950 L/ha) of water.	
	Root Dip	Mix a 0.5% v/v solution (2 quarts/100 gallons of water or 5 L/1000 L of water).	Apply as pre-plant dip to the roots for 30 seconds. Plant within 48 hours after dipping. Mix a fresh solution daily.
	Chemigation		
	Overhead:	Apply 2-3 quarts/acre (5-7 L/ha) in a minimum of 1,000 gallons/acre (9,350 L/ha) of water.	Apply with normal irrigation schedule. Follow Disease Programs* as stated above. Do not apply more than 6 times per crop cycle.
	Low Volume:	Apply 2-4 quarts/acre (5-9 L/ha) in a minimum of 100 gallons/acre (950 L/ha) of water.	

* Check with your local Extension Agent or Crop Consultant if you are unsure about disease prevention, control or severity/pressure.

** See the label booklet for Fungi-Phite for a full list of diseases pertaining to each this crop group.

† Fungi-Phite has suppressive properties on diseases caused by certain bacteria or fungi. When applying Fungi-Phite for disease suppression, use it in combination with another registered bactericide or fungicide registered for the same crops listed on the label.

Read the "Conditions of Sale and Limitations of Warranty and Liability" in the label booklet for Fungi-Phite before using.