Supplemental Label



GROUP 7D INSECTICIDE

DISTANCE® Insect Growth Regulator EPA Reg. No. 59639-96 (Except California and New York)

DISTANCE INSECT GROWTH REGULATOR USE ON: INDOOR GROWN FRUITING VEGETABLES

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

RESISTANCE MANAGEMENT RECOMMENDATIONS

For resistance management purposes, *Distance* Insect Growth Regulator is a Group 7D insecticide. Any insect population may contain individuals naturally resistant to *Distance* and other Group 7D insecticides. The resistant individuals dominate the insect population if these insecticides are used repeatedly. These resistant insects may not be controlled by *Distance* Insect Growth Regulator or other Group 7D insecticides, although local experts should be consulted for local resistance recommendations. The classification scheme is based on mode of action. It is recognized that resistance of insects and mites to insecticides and acaricides can also result from enhanced metabolism, reduced penetration or behavioral changes that are not linked to any site of action classification but are specific for individual chemicals or chemical groupings. Despite this, alternation of compounds from different chemical classes remains a viable management technique.

To delay insecticide resistance

- Avoid exclusive repeated use of insecticides from the same chemical subgroup.
- Integrate other control methods (chemical, cultural, biological) into insect control programs.

For further information contact your local Pest Control Advisor (PCA).

Table 1. Directions for Use on Indoor Grown Fruiting Vegetables.

PESTS	RATES	APPLICATION METHOD	SPECIAL INSTRUCTIONS		
Aphids (suppression) Greenhouse	6 fl. oz./ 100 gals.	Foliar Spray 100 gals. of spray mix will	Apply the spray mixture uniformly to all plant surfaces and to the point of runoff. Make first application when adult insects begin to appear. If necessary, make a second application from 14 to 28 days after the first application. If an additional application is needed fewer than 14 days after the first treatment, use an IGR with another mode of action or another chemical class of insecticide. Use lower rate and longer interval for newly established infestations and when plants are not rapidly flushing new growth. Use higher rates and shorter interval for established infestations and/or when plants are rapidly flushing new growth. Apply no more than two times per cropping cycle or no more than two times per six months. If rapid control of adult insects is required, apply a labeled adulticide.		
Whitefly Silverleaf Whitefly Sweetpotato Whitefly		treat 20,000 sq. ft. of area.			
Fungus Gnats Shore Flies	3 to 6 fl. oz./100 gals.	Sprench 100 gals. of spray mix will treat 5,000 sq. ft. of area.	For the control of fungus gnats and shore flies apply to potting media as a heavy, coarse spray (sprench) through conventional equipment to all insect infested surfaces or where insects may breed. Complete coverage of infested areas is essential for control. For optimal control, treat breeding areas under benches at the same time that the crop is treated. The soil surface should be moist at the time of application.		
	Broadcast Application to Soil Surface: For bed, bench and container grown plants, apply Distance coarse spray or sprench to the soil surface. Mix 3 to 6 fl. Distance in 100 gals. of water and apply to the soil surface volume of 2 to 3 gals. of final spray solution per 100 sq. ft. o If a second application of Distance is needed, allow a minim 21 days between applications. Apply no more than two time cropping cycle or no more than two times per six months.				e. Mix 3 to 6 fl. oz. of the soil surface at a per 100 sq. ft. of area. d, allow a minimum of the than two times per er six months.
Fungus Gnats Shore Flies 2 fl. oz./ 100 gals. Saturate only the top 1" to 1.5" of soil. For the control of fungus gnats and shore media as a drench application through co For optimal control, apply additional amount breeding areas under benches at the same treated. The soil surface should be moist at				ventional equipment. ts of spray solution to time that the crop is	
	Drench Application to Soil Surface of Individual Mix 2 fl. oz. of Distance in 100 gals. of water and essurface of potting media to ensure uniform treatment oz. of finished solution per 6 inch pot. Adjust volunt for smaller or larger pots (see drench mixing chart to saturate potting media with drench solution, on 1.5" of soil needs to be drenched in order to ack control. Do not drench plants more than one time process.				er and evenly apply to treatment. Apply 3 fl. st volume accordingly chart below). Do not ion , only the top 1" to to achieve effective e time per crop cycle.
			Pot Diameter (inches)	Drench Volume (fl. oz./pot)	Rate/100 Gals. (fl. oz.)
			4	1	2
			5	2	2
			6	3	2
			8	5	2
			10 12	7 10	2 2
			IΖ	10	

NOTE: Since fruiting vegetable varieties are numerous, constantly changing and may react differently to DISTANCE and tank mixtures including *Distance*, test the product(s) on a small scale before making large-scale applications.

Do not apply to tomato varieties less than 1 inch in diameter.

Do not apply to non-bell peppers.

Do not apply within one (1) day of harvest.

Do not make more than two (2) Distance applications per season.

Do not exceed 12 fl. oz. Distance per acre per season.

Regardless of formulation, do not apply more than 0.176 lb. ai of pyriproxyfen per acre per season.

THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF APPLICATION. PLEASE REFER TO CONTAINER LABEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS. FOLLOW ALL APPLICATION DIRECTIONS, RESTRICTIONS, AND PRECAUTIONS ON THE EPA REGISTERED LABEL.

PLEASE CONTACT VALENT U.S.A. CORPORATION AT 1-800-6-VALENT (682-5368) TO DETERMINE IF THIS USE IS REGISTERED IN YOUR STATE.

Manufactured for Valent U.S.A. Corporation P. O. Box 8025 Walnut Creek, CA 94596-8025

Made in U.S.A.



Products That Work, From People Who Care™ | www.valent.com | 800-89-VALENT (898-2536) Read and follow the label instructions before using.

Distance is a registered trademark and Products That Work, From People Who Care is a trademark of Valent U.S.A. Corporation.©2006 Valent U.S.A. Corporation. All rights reserved. Printed in the USA. 2006-DIS-0010 Ex CA & NY 3/06 mf/AV