

PYRAFLUFEN-ETHYL GROUP 14 HERBICIDE

# EDICT™ 4SC

IVM HERBICIDE

**ACTIVE INGREDIENT:**

**Pyraflufen-ethyl:** Acetic acid, [2-chloro-5-[4-chloro-5-(difluoromethoxy)-1-methyl-1H-pyrazol-3-yl]-4-fluorophenoxy]-, ethyl ester ..... 4.0%

**OTHER INGREDIENTS:** ..... 96.0%

**TOTAL** ..... 100.0%

Contains 0.35 lb pyraflufen-ethyl per gallon

(AYE)

EPA Reg. No. 71711-69

EPA Est. No. 92534-NC-1 92534-TX-2 54675-MEX-001

superscript corresponds to lot number

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

*See inside booklet for First Aid, Precautionary Statements, and Directions for Use*

**NET CONTENTS: 250 gallons**

 **NICHINO**

500521 01/25

## FIRST AID

### If on skin or clothing

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

## HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor or going for treatment. For additional information on this pesticide product, including human health concerns and medical emergencies, call 1-800-348-5832. In case of fire or spills, information may be obtained by calling 1-800-424-9300.

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

## CAUTION

Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

### PERSONAL PROTECTIVE EQUIPMENT (PPE)

#### Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves made of barrier laminate, butyl rubber  $\geq 14$  mils, nitrile rubber  $\geq 14$  mils, neoprene rubber  $\geq 14$  mils, natural rubber  $\geq 14$  mils, polyethylene, polyvinyl chloride (PVC)  $\geq 14$  mils, and/or Viton™  $\geq 14$  mils

#### User Safety Requirements

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

## Engineering Controls

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

### User Safety Recommendations

Users should:

- Wear long sleeved shirt and long pants, shoes plus socks, and chemical and/or water-resistant gloves.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

## ENVIRONMENTAL HAZARDS

This product is toxic to fish and aquatic invertebrates. This product may contaminate water through drift of spray in wind or via runoff events. Use care when applying in areas adjacent to any body of water. **DO NOT** apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment washwater or rinsate. **DO NOT** apply when weather conditions favor drift from treated areas.

### Nontarget Organism Advisory

This product is toxic to plants and may adversely impact the forage and habitat of nontarget organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of nontarget organisms by following label directions intended to minimize spray drift.

## DIRECTIONS FOR USE

**It is a violation of federal law to use this product in a manner inconsistent with its labeling.**

**DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the state or tribal agency responsible for pesticide regulation.

## NONAGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. For other uses, including interiorscapes and other nonagricultural uses, **DO NOT** enter treated areas without protective clothing until sprays have dried.

## PRODUCT INFORMATION

**EDICT™** 4SC IVM Herbicide is a contact herbicide and requires thorough coverage for complete broadleaf weed control. **EDICT** 4SC IVM Herbicide must be tank mixed with another foliar active broadleaf herbicide for complete control of most broadleaf weeds.

**DO NOT** apply **EDICT** 4SC IVM Herbicide through any type of irrigation system.

**EDICT** 4SC IVM Herbicide is rainfast one hour after application.

## WEEDS CONTROLLED

The following broadleaf weed species can be controlled or suppressed up to 4 inches in height or less or rosettes of 3 inches in diameter or less. Tank mixtures of **EDICT** 4SC IVM Herbicide with other labeled broadleaf herbicides may be needed for control of some weed species. Control may be reduced with weeds larger than 4 inches in height or 3 inches in diameter.

Alkaliweed\*  
Amaranth, Palmer\*  
Bedstraw  
Beggartick, hairy  
Beggarweed, Florida  
Bindweed, field  
Buckwheat, wild  
Canola  
Carpetweed  
Celery, wild  
Chickweed  
Clover, white  
Cocklebur

Cotton, volunteer  
(conventional, GMO varieties)  
Dandelion, common  
Dock, curly  
Dollarweed  
Eclipta  
Eveningprimrose, cutleaf  
Fleabane\*  
Geranium, Carolina  
Henbit  
Horsenettle\*  
Kochia  
Ladysthumb

Lambsquarters, common  
Lettuce, prickly  
Mallow, common  
Malva spp.  
Marestail\*  
Milkthistle  
Morningglory species  
Mustard, wild\*  
Nettle, stinging  
Nightshade, black  
Nightshade, silverleaf  
Panicle willowweed  
Pigweed, redroot

**K 998 G7 CBHFC @98** (continued)

Pigweed, smooth Pineapple-weed Poinsettia, wild Poison-ivy Potato, volunteer Prickly sida (teaweed) Purslane, common Radish, wild Ragweed, common	Ragweed, giant Redmaid Rocket, London Sesbania, hemp Smartweed, Pennsylvania Smellmelon Sowthistle, annual Spurge, leafy Sunflower, common	Tansymustard, Western Thistle, Canada Thistle, Russian Toadflax, Dalmatian Velvetleaf Virginia-creeper Waterhemp, common Waterhemp, tall
*suppression		

**K 998 F9G-GH5 B79 A5 B5 ; 9A9BH**

For resistance management, **987 H4SC** IVM Herbicide is a Group 14 herbicide. Any weed population may contain or develop plants naturally resistant to **987 H4SC** IVM Herbicide and other Group 14 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Appropriate resistance management strategies must be followed.

Proactively implement diversified weed control strategies to minimize selection for weed populations resistant to one or more herbicides. A diversified weed management program may include the use of multiple herbicides with different modes of action and overlapping weed spectrum with or without tillage operations and/or other cultural practices. Research has demonstrated that using the labeled rate and application instructions is important to delay the selection for resistance. Scouting after a herbicide application is important because it can facilitate the early identification of weed shifts and/or weed resistance and thus provide direction on future weed management practices. One of the best ways to contain resistant populations is to implement measures to avoid allowing weeds to reproduce by seed or to proliferate vegetatively. Cleaning equipment between sites and avoiding movement of plant material between sites will greatly aid in retarding the spread of resistant weed seed.

To delay herbicide resistance, take one or more of the following steps:

- Rotate the use of **987 H4SC** IVM Herbicide or other Group 14 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds in a field.

- Use tank mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner. Consult your local extension service or certified crop advisor if you are unsure as to which active ingredient is currently less prone to resistance.
- Adopt an integrated weed management program for herbicide use that includes scouting and uses historical information related to herbicide use and that considers tillage (or other mechanical control methods), cultural (e.g., higher crop seeding rates; precision fertilizer application method and timing to favor the crop and not the weeds), biological (weed-competitive crops or varieties), and other management practices.
- Scout after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of noncontrolled plants of a particular weed species; (3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method (for example, hoeing or tillage). Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields and planting clean seed.
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your local extension specialist or certified crop advisor for additional pesticide resistance management and/or integrated weed management recommendations for specific crops and weed biotypes.
- Report lack of performance to registrant or their representative.

Suspected herbicide resistant weeds may be identified by these indicators:

- Failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds;
- A spreading patch of noncontrolled plants of a particular weed species; and
- Surviving plants mixed with controlled individuals of the same species.

Contact your local sales representative, certified crop advisor or pest control advisor, or extension agent to find out if suspected resistant weeds to this MOA have been found in your region. If resistant biotypes of target weeds have been reported, use the application rates of this product specified for your local conditions. Tank mix products so that there are multiple effective mechanisms of action for each target weed.

## **MANDATORY SPRAY DRIFT MANAGEMENT**

### **Ground Boom Applications**

- User must only apply with the release height specified by the manufacturer but no more than 3 feet above the ground or crop canopy.
- Applicators are required to use fine or coarser droplet size (ASABE S572.1).
- **DO NOT** apply when wind speeds exceed 10 miles per hour at the application site.
- **DO NOT** apply during temperature inversions.

### **Boomless Ground Applications**

- Applicators are required to use fine or coarser droplet size (ASABE S572.1) for all applications.
- **DO NOT** apply when wind speeds exceed 10 miles per hour at the application site.
- **DO NOT** apply during temperature inversions.

## **SPRAY DRIFT ADVISORIES**

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.

BE AWARE OF NEARBY NONTARGET SITES AND ENVIRONMENTAL CONDITIONS.

### **Importance of Droplet Size**

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

### **Controlling Droplet Size - Ground Boom**

**Volume:** Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.

**Pressure:** Use the lowest spray pressure directed for the nozzle to produce the target spray volume and droplet size.

**Spray Nozzle:** Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

### **Boom Height - Ground Boom**

Use the lowest boom height that is compatible with the spray nozzles that will provide uniform coverage. For ground equipment, the boom must remain level with the crop and have minimal bounce.

### **Shielded Sprayers**

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

### **Temperature and Humidity**

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

### **Temperature Inversions**

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.



## **Wind**

Drift potential increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS. Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

## **Boomless Ground Applications**

Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.

## **Handheld Technology Applications**

Take precautions to minimize spray drift.

## **TANK MIXTURES**

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and application instructions on all product labels involved in tank mixing. Users must follow the most restrictive application instructions and precautionary statements of each product in the tank mixture.

**EDICT** 4SC IVM Herbicide may be applied as a tankmix or in sequential application with other herbicide, fungicide, or insecticide products. Weather, crop conditions, or the presence of certain weeds, crop damaging insects, or diseases will indicate the inclusion of other pesticides in the application.

Note: Test compatibility of **EDICT** 4SC IVM Herbicide in any tankmix combination before use. To determine the physical compatibility with other products, use a jar test, as described below:

Using a quart jar, add the proportionate amounts of the products to 1 quart of water. Add wettable powders and water-dispersible granular products first, then liquid flowables, and emulsifiable concentrates last. After thoroughly mixing, let stand for at least 5 minutes. If the combination remains mixed or can be remixed readily, it is physically compatible. Once compatibility has been proven, use the same procedure for adding required ingredients to the spray tank.

FYUX'UbX'Z`ck`U`~`UVY`X'fYV'cbg'Z'f`YUW`HUb\_a`Jl`dfcXi`W' 5`k`Ung'i`gY`b`UW'fXUbW'k`Jh`H`Y'a`cghfYghf`W'j`Y'cZ`UVY`dfYU' h'cbg'UbX`"ja`Jh'cbg"

## A-L-B; `8-F97H-CBG

**98-7 H' (G7`-JA`<YfV'WXY`5`cbY.'** Fill spray tank with  $\frac{3}{4}$  of the amount of water needed for the intended application, and then turn on agitation. Pour the specified amount of product on the surface of the water in the spray tank. Add the remaining water volume to the spray tank with agitation running. Keep agitation running during filling and spraying operations. If spraying must be stopped before emptying the sprayer, resume agitation before spraying the remainder of the load. Mix only as much spray solution as can be sprayed within four hours. Storage and use of the previous day's spray mix may result in reduced activity.

**98-7 H' (G7`-JA`<YfV'WXY`Jb`HUb`\_A`Jl`h`fYg.'** Begin with clean equipment. Fill spray tank with  $\frac{3}{4}$  of the amount of water needed for the intended application, and turn on agitation. If using a buffering agent, add after filling the tank with  $\frac{3}{4}$  amount of water. Add the specified amount of tankmix products in the following order while maintaining agitation:

- 1) products in water-soluble packets
- 2) wettable powders
- 3) water-dispersible granulars and/or soluble powders
- 4) flowable liquids (including **98-7 H'4SC IVM Herbicide**)
- 5) emulsifiable concentrates
- 6) adjuvants and/or oils
- 7) remaining amount of water to achieve the desired level

Always follow the labeled mixing instructions of any partner products. Keep agitation running during filling and spraying operations. If spraying must be stopped before emptying the sprayer, resume agitation before spraying the remainder of the load. Mix only as much spray solution as can be sprayed within four hours. Storage and use of the previous day's spray mix may result in reduced activity.

Use an approved agricultural buffering agent, buffering to pH 7.5 or less if using **98-7 H'4SC IVM Herbicide** in a water source greater than or equal to pH 7.5. Always buffer the water source BEFORE adding **98-7 H'4SC IVM Herbicide** to the spray tank.

## EQUIPMENT CLEANING

**DO NOT** allow the spray solution to dry in the application equipment. After application and before using the sprayer equipment for any other applications, the sprayer must be thoroughly cleaned. Applicators must ensure proper equipment clean-out for any other products mixed with **EDICT 4SC IVM Herbicide** as provided on the other product label(s). Immediately following application, clean all equipment thoroughly with detergent or a spray tank cleaner and water as described below. Residues of **EDICT 4SC IVM Herbicide** remaining in inadequately cleaned equipment and released in subsequent applications can cause injury to crops.

1. Drain sprayer tank, hoses, and spray boom, and thoroughly rinse with clean water the inside of the spray tank, sprayer hoses, boom, and nozzles to remove any sediment or residues.
2. Fill the tank  $\frac{1}{2}$  full with clean water, add the appropriate detergent (follow manufacturer's directions for use). Fill tank to capacity, and operate the sprayer with agitation for 15 minutes to flush hoses, boom, and nozzles.
3. Drain the sprayer tank, lines, and booms. Rinse the tank with clean water and flush through the hoses, boom, and nozzles. Remove and clean spray nozzles, tips, and screens.
4. Dispose of all cleaning solutions, rinsate, and washwaters in accordance with federal, state, and local regulations.

## SPOT TREATMENT

For spot treatment to listed broadleaf weeds, refer to the information below to determine the amount of **EDICT 4SC IVM Herbicide** to add to a tank. Spray using a pressure (pump-up) sprayer (or similar application equipment) until wet but prior to runoff. Use information for rates, concentrations, water volumes, and timing and frequency of application can be found in the **Rate/Acre** and **Application Instructions** columns in the **Use Directions** tables. Please refer to and follow all precautions and restrictions under **Application Instructions** for the crop to be treated.

### Fluid Ounces of EDICT 4SC IVM Herbicide to Add to Sprayer Tank

Sprayer tank capacity (gallons)	Spray volume (gallons/A)	fl oz EDICT 4SC IVM Herbicide to add to spray tank for an application rate of	
		0.5 fl oz/A (0.0013 lb ai/A)	1.9 fl oz/A (0.0053 lb ai/A)
1	20	0.03	0.10
3	20	0.08	0.29
5	20	0.13	0.48
10	20	0.25	0.95

#### Formula

Fluid Ounce **EDICT** 4SC IVM Herbicide to add to sprayer tank = 
$$\frac{\text{Application Rate} \times \text{Sprayer Tank Capacity}}{\text{Spray Volume}}$$

*(continued)*

### Fluid Ounces of EDICT 4SC IVM Herbicide to Add to Sprayer Tank *(continued)*

#### Example Calculation for 1 gallon sprayer tank capacity

Fluid Ounce **EDICT** 4SC IVM Herbicide to add to sprayer tank =

$$\frac{1.9 \text{ fl oz/A} \times 1 \text{ gallon}}{20 \text{ gallons/A}} = 0.10 \text{ fl oz}$$

$$20 \text{ gallons/A} = 0.10 \text{ fl oz}$$

where: Application rate = 1.9 fl oz/A

Sprayer tank capacity = 1 gallon

Spray volume = 20 gallons/A

#### Example Calculation for 5 gallon sprayer tank capacity

Fluid Ounce **EDICT** 4SC IVM Herbicide to add to sprayer tank =

$$\frac{1.9 \text{ fl oz/A} \times 5 \text{ gallons}}{20 \text{ gallons/A}} = 0.48 \text{ fl oz}$$

$$20 \text{ gallons/A} = 0.48 \text{ fl oz}$$

where: Application rate = 1.9 fl oz/A

Sprayer tank capacity = 5 gallons

Spray volume = 20 gallons/A

## USE DIRECTIONS

### Noncrop Weed Control

airports and airfields; commercial plants; storage and lumber yards; fence lines and fence rows; farmyards and farm buildings; barrier strips and firebreaks; equipment areas; railroads; roadside and utility rights-of-way; fuel tank farms and pumping stations; dry ditches and ditchbanks; vacant lots; and similar agricultural and industrial noncrop sites

Pest	Rate/Acre	Application Instructions
Listed Broadleaf Weeds	1.5 to 1.9 fl oz/acre (0.0040 to 0.0053 lb ai/acre)	<ul style="list-style-type: none"><li>• Apply in a minimum of 20 to 40 gallons spray solution per acre by ground.</li><li>• Avoid contact with desirable vegetation.</li><li>• Use of a COC adjuvant at a concentration of 1% to 2% is directed for optimum weed control. Use the higher COC rate for larger labeled weed species or in low moisture conditions.</li><li>• COC adjuvants are advised, although other adjuvants may be used.</li><li>• Use the higher rate for hard-to-control weeds.</li></ul>

### USE RESTRICTIONS

- **DO NOT** exceed 1.9 fl oz (0.0053 lb ai) of **EDICT 4SC IVM** Herbicide per acre in a single application for this use.
- **DO NOT** make more than 3 applications or exceed 5.8 fl oz (0.0159 lb ai) of **EDICT 4SC IVM** Herbicide per acre per year.
- Allow a minimum of 30 days between applications for this use.

## STORAGE AND DISPOSAL

**DO NOT** contaminate water, food, or feed by storage or disposal.

**STORAGE:** Store in original container, and keep tightly closed when not in use. Store in a cool, dry place.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

### CONTAINER HANDLING:

**Nonrefillable plastic container (less than 5 gallons).** Nonrefillable container. **DO NOT** reuse or refill this container. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container  $\frac{1}{4}$  full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available, or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill or by other procedures approved by state and local authorities.

**Nonrefillable plastic container (greater than 5 gallons).** Nonrefillable container. **DO NOT** reuse or refill this container. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container  $\frac{1}{4}$  full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or a mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available, or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill or by other procedures approved by state and local authorities.

## IMPORTANT: READ BEFORE USE

By using this product, user or buyer accepts the following conditions, warranty, disclaimer of warranties, and limitations of liability.

**CONDITIONS:** The directions for use of this product are believed to be accurate and must be followed carefully. However, because of extreme weather and soil conditions, use methods and other factors beyond the control of Nichino America, Inc. (NAI), it is impossible for NAI to eliminate all risks associated with the use of this product. As a result, crop injury or ineffectiveness is always possible. To the extent consistent with applicable law, all such risks are assumed by the user or buyer.

**DISCLAIMER OF WARRANTIES:** TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THERE ARE NO WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, WHICH EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of NAI is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, NAI disclaims any liability whatsoever for incidental or consequential damages, including, but not limited to, liability arising out of breach of contract, express or implied warranty (including warranties of merchantability and fitness for a particular purpose), tort, negligence, strict liability, or otherwise.

**LIMITATIONS OF LIABILITY:** TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES, OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY, OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT THE ELECTION OF NICHINO AMERICA, INC., THE REPLACEMENT OF PRODUCT.

©2025 Nichino America, Inc. Edict is a trademark of Nichino America, Inc. Viton is a trademark of The Chemours Company.

Formulated and Packaged for  
Nichino America, Inc.  
4550 Linden Hill Road, Suite 501  
Wilmington, DE 19808  
888-740-7700

71711-69.20230504.1  
06/02/2023



# EDICT™ 4SC

IVM HERBICIDE

## ACTIVE INGREDIENT:

**Pyraflufen-ethyl:** Acetic acid, [2-chloro-5-[4-chloro-5-(difluoromethoxy)-1-methyl-1H-pyrazol-3-yl]-

4-fluorophenoxy]-, ethyl ester ..... 4.0%

**OTHER INGREDIENTS:** ..... 96.0%

**TOTAL** ..... 100.0%

Contains 0.35 lb pyraflufen-ethyl per gallon

(AYE)

EPA Reg. No. 71711- 69 EPA Est. No. 92534-NC-1 92534-TX-2 54675-MEX-001  
superscript corresponds to lot number

## KEEP OUT OF REACH OF CHILDREN CAUTION

### FIRST AID

If on skin  
or clothing

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

### HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor or going for treatment. For additional information on this pesticide product, including human health concerns and medical emergencies, call 1-800-348-5832. In case of fire or spills, information may be obtained by calling 1-800-424-9300.

PYRAFLUFEN-ETHYL

GROUP

14

HERBICIDE

## STORAGE AND DISPOSAL

**DO NOT** contaminate water, food, or feed by storage or disposal.

**STORAGE:** Store in original container, and keep tightly closed when not in use. Store in a cool, dry place. **PESTICIDE DISPOSAL:**

Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility. **CONTAINER**

**HANDLING:** Nonrefillable container. DO NOT reuse or refill this container. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or a mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available, or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill or by other procedures approved by state and local authorities.

**See inside booklet for First Aid, Precautionary Statements, and Directions for Use**

Formulated and Packaged for

Nichino America, Inc.

4550 Linden Hill Road

Suite 501

Wilmington, DE 19808

888-740-7700

**NET CONTENTS: 250 gallons**

500521 01/25