

Product Bulletin



Loveland Products, Inc., P.O. Box 1286, Greeley, Colorado 80634

Reaper® ClearForm™

EPA Reg. No. 34704-1078

2(ee) Recommendation[†]

Control of Additional Pest in Florida

ATTENTION

- [†]This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA or state lead pesticide agency.
- Read and follow all applicable directions for use, precautions and limitations on the federally registered label attached to the product container.

Additional pest on avocados controlled within existing rate range of the federally registered product label for Reaper ClearForm:

Avocados	
Additional Target Pest	Instructions
Mirids (<i>Daghbertus fasciatus</i> , <i>Daghbertus olivaceous</i> , <i>Rhinacloa</i> sp.)	Apply 10-20 fl. oz. Reaper ClearForm per acre. For best results, apply when immature mirids are first observed. Apply in the January to April time period. See restrictions below.

Restrictions:

- The user must refer to the federally approved **Reaper ClearForm** label and read and follow all applicable directions for use, restrictions, and precautions.
- Do not apply within 14 days of harvest.
- Ground and aerial application permitted
- Wait at least 30 days before repeating application
- Do not make more than 2 applications of Reaper ClearForm or any other foliarly-applied abamectin-containing product per season.
- Do not apply more than 40 fl. oz. per acre (or 0.047 pounds active ingredient per acre) of Reaper ClearForm or any other foliarly applied abamectin-containing product per season.
- Do not apply more than 20 fl. oz. per acre (or 0.023 pounds active ingredient per acre) of Reaper ClearForm or any other foliarly applied abamectin-containing product per application.

- Do not allow livestock to graze in treated orchards.

The user must refer to the federally approved Reaper® ClearForm™ label and read and follow all applicable directions for use, restrictions, and precautions.

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

This recommendation should be in the possession of the user at the time of pesticide application.