

TECHNICAL INFORMATION BULLETIN

SPINOSAD

GROUP

5

INSECTICIDE



EPA Reg. No. 91234-279

FIFRA Section 2(ee) Recommendation

This recommendation is made as permitted by Section 2(ee) of FIFRA and has not been submitted to or approved by the U.S. Environmental Protection Agency.

**FOR DISTRIBUTION AND USE ONLY WITHIN THE STATES OF IDAHO, MINNESOTA,
NORTH DAKOTA, OREGON AND WASHINGTON**

CONTROL OF POTATO PSYLLID

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product inconsistent with its labeling.
- This product bulletin must be in the possession of the user at the time of pesticide application.
- Read the "Limitation of Warranty and Liability" in the label booklet for **Falcondor SC** before using.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

Refer to product label for **Falcondor SC** for General Use Precautions, Mixing Directions and Application Directions.

Apply 5.04 fluid ounces of **Falcondor SC** per acre to potatoes for control of potato psyllid.

Treat when pests appear, targeting eggs at hatch or small larvae. When plants are growing rapidly, repeat applications may be necessary to protect new foliage.

Preharvest Interval: Do not apply within 7 days of harvest.

Minimum Treatment Interval: Do not make applications less than 7 days apart.

Follow resistance management recommendations on the product label.

Consult your local Atticus, LLC representative, extension service specialist, certified crop advisor, or state agricultural experiment station for any additional local use recommendations for your area.



Manufactured for:

Atticus, LLC

940 NW Cary Parkway, Suite 200
Cary, NC 27513