

THIS RECOMMENDATION IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE EPA.

## 2(ee) Recommendation



IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING

This Product Bulletin Contains New or Additional Directions For Use Which Are Recommended By Sym-Agro, Inc. Under FIFRA Section 2(ee)

FOR DISTRIBUTION AND USE ONLY IN  
Arizona, California, Oregon, Washington, Idaho, Montana, Utah

EPA REG. No. 84876-2

| CROP                           | PEST  | RATE OF APPLICATION                          |
|--------------------------------|---|--|
| Stone Fruit<br>(Crop Group 12) | Alternaria, Alternaria late blight ( <i>Alternaria alternata</i> )<br>Leaf spots ( <i>Alternaria</i> spp.)<br>Blue mold ( <i>Penicillium</i> spp.)<br>Gray Mold Botrytis blossom blight ( <i>Botrytis cinerea</i> )<br>Powdery mildew ( <i>Oidium</i> spp., <i>Podosphaera leucotricha</i> , <i>Sphaerotheca pannosa</i> )<br>Ripe fruit rot ( <i>Monilinia fruticola</i> , <i>Monilinia laxa</i> , <i>Botrytis cinerea</i> , <i>Rhizopus</i> spp.)<br>Peach Leaf Curl ( <i>Taphrina deformans</i> )<br>Rust ( <i>Tranzschelia discolor</i> )<br>Scab ( <i>Venturia carpophila</i> )<br>Shot Hole ( <i>Wilsonomyces carpophilus</i> ) | 20.4 – 45.7 fl oz/A<br>(0.34 – 0.75 lb ai/A) |

### DIRECTIONS FOR USE:

**Botrytis:** For control of Botrytis gray mold apply ProBlad® Verde prior to onset of disease development when conditions favor Botrytis development up to the harvest.

**Powdery mildew:** Begin applications of ProBlad® Verde preventively when conditions are favorable for disease development and continue on a 7 to 10-day interval if conditions continue to favor disease development. Under conditions of severe disease pressure, use the higher labeled rate.

**General:** Increase spray volume as growth increases in order to provide thorough coverage of foliage and fruit for optimum disease control.

### RESTRICTIONS:

Do not make more than two sequential applications of ProBlad® Verde Fungicide before alternating to a labeled fungicide with a different mode of action.

Do not make more than 5 applications of ProBlad® Verde Fungicide per harvest cycle.

**PHI: 1 Day**

**ALL APPLICABLE DIRECTIONS, RESTRICTIONS, WORKER PROTECTION STANDARDS REQUIREMENTS AND PRECAUTIONS ON THE EPA REGISTERED PROBLAD VERDE® FUNGICIDE (EPA REG. No. 84876-2) LABEL MUST BE FOLLOWED. THIS RECOMMENDATION MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.**

ProBlad is a trademark of CEV S.A.  
ProBlad Verde\_2ee\_Stone Fruit Crop Expanded Diseases\_05/05/2022



Sym-Agro, Inc.  
PO Box 71, Visalia, CA 93291