

FIFRA 2(ee) Recommendation

EPA Reg. No. 279-9629

FOR DISTRIBUTION AND USE IN REGISTERED STATES (EXCEPT NEW YORK AND TEXAS)

THIS RECOMMENDATION, WHICH CONTAINS ADDITIONAL DIRECTIONS FOR USE, IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE US EPA.

This recommendation for AVAUNT® eVo insect control is valid until June 30, 2030, or until withdrawn, canceled or suspended.

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL MUST BE FOLLOWED.

THESE USE DIRECTIONS MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

FOR SUPPRESSION OF ADULT PEPPER WEEVIL IN PEPPER

Directions For Use

CROP	APPLICATION METHOD	TARGET PEST	RATE	PHI (Days)	REI (Hours)
Pepper (BELL, CHILI, COOKING, PIMENTO AND SWEET)	Foliar	Pepper weevil (Anthonomus eugenii)	6.0 oz/A (0.11 lb ai/A)	3	12

- AVAUNT eVo insect control is recommended as a foliar application for suppression of adult pepper weevil.
- Apply AVAUNT eVo insect control at 6.0 oz/A (0.11 lb ai/A) as part of a foliar program rotating with different mode of action insecticides known to be effective on pepper weevil.
- Spray programs should begin at flowering or at the first sign of adult weevils.
- Make repeat applications on a 5 to 7-day spray schedule if monitoring indicates the presence of adult weevils.
- See the Resistance Management section of the AVAUNT eVo insect control Section 3 Federal label for additional guidance regarding insecticide resistance management.



 Thorough coverage is essential to achieve best results. Select a spray volume appropriate for the size of plants and density of foliage being treated.

RESTRICTIONS AND PRECAUTIONS

- Do not apply more than 24 oz/A AVAUNT eVo insect control or 0.45 lb ai of indoxacarb-containing products per acre per crop.
- Do not make more than 4 applications per crop.
- Do not apply more than 72 oz/A AVAUNT eVo insect control or 1.35 lb ai of indoxacarb-containing products per acre per year.
- Apply in a minimum of 10 gallons of water per acre (GPA) by ground.
- The minimum interval between sprays is 5 days.

IMPORTANT NOTICE

BEFORE BUYING OR USING FMC PRODUCT(S) COVERED BY THIS BULLETIN, READ THIS BULLETIN, AND THE LABELS FOR ALL PRODUCTS REFERENCED IN THIS BULLETIN, IN THEIR ENTIRETY, AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS.

It is a violation of Federal law to use this or any other pesticide product in a manner inconsistent with its labeling. This Bulletin contains supplemental instructions for use of the referenced FMC product. Follow these instructions carefully. This Bulletin must be in the possession of the User at the time of application. Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using the FMC product(s) covered by this Bulletin. FMC will not be responsible for losses or damages resulting from use of this product in any manner not specifically stated by FMC on the Section 3 label, or other labels, or this Bulletin. User assumes all risks associated with any such nonspecified use. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability of the Section 3 Federal product label.

R-4971 060425 06-04-25 (Replaces R-4440 033020 03-26-20)

FMC, the FMC logo and Avaunt are trademarks of FMC Corporation or an affiliate. ©2025 FMC Corporation

