

TECHNICAL INFORMATION BULLETIN

GROUP 15 INSECTICIDE

RIMON[®] 0.83EC Insecticide

EPA Reg. No. 66222-35-400

FIFRA Section 2(ee) Recommendation

This recommendation is made as permitted by Section 2(ee) of FIFRA and has not been submitted to or approved by the U.S. Environmental Protection Agency.

FOR USE AND DISTRIBUTION IN THE STATE OF CALIFORNIA ONLY

**The addition of THRIPS control in
Highbush Blueberry**

DIRECTIONS FOR USE

Use information in accordance with Section 2(ee) of FIFRA.

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Read and follow all applicable directions for use, restrictions, worker protection standard requirements, and precautions on the EPA registered main label attached to the container of **RIMON[®] 0.83EC Insecticide (EPA Reg. No. 66222-35-400)**.

022817V024CA2ee

BLUEBERRY (Highbush)

Target Pest	Rate (fl oz / A)	Application Instructions
Thrips	20 - 30	<p>Make application when adults are observed and prior to egg laying.</p> <p>Applications may also be made when the majority of the population is at egg hatch to the second instar.</p> <p>Transovarial control via females laying non-viable eggs may occur if applications are made when adults are first observed and prior to egg laying.</p>
<ul style="list-style-type: none">• Some phytotoxic symptoms to foliage in the form of mottled chlorosis may be observed when RIMON 0.83EC Insecticide is applied to blueberries under conditions of high temperatures and/or drought stress, particularly during periods of new tender shoot growth. Such phytotoxic symptoms will not occur on future growth and will not affect fruiting or yields. Higher spray volumes and lower spray concentration will minimize the risk of transient phytotoxic symptoms on newly expanded foliage.• Use higher rates and higher spray volumes when larvae are large or foliage canopy is tall or dense.• Repeat applications as needed to protect new foliage growth and fruit but do not apply less than 10 days apart.• Do not apply more than 90 fl oz per acre per year (0.58 lb ai per acre per year).• Do not apply within 8 days of harvest.		

Distributed By:

MacDermid Agricultural Solutions, Inc.
c/o ARYSTA LIFESCIENCE NORTH AMERICA, LLC
15401 Weston Parkway, Suite 150
Cary, NC 27513

Arysta LifeScience and the Arysta LifeScience logo are registered trademarks of Arysta LifeScience Corporation.

RIMON is a registered trademark of ADAMA Makhteshim Ltd.

©2018 Arysta LifeScience Group Company.