



FIFRA Section 24(c)
SPECIAL LOCAL NEED LABEL

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF **ARIZONA**

ASSAIL® 30SC INSECTICIDE

FOR USE ON COTTON AT A HIGHER RATE

This label is valid until December 31, 2027 or until otherwise amended, withdrawn, cancelled or suspended.

ACTIVE INGREDIENT: acetamiprid (E)- N¹-[(6-chloro-3-pyridyl)methyl] – N²-cyano-N¹-methyl acetamidine..... 30.0% by wt.
OTHER INGREDIENTS..... 70.0% by wt.
TOTAL.....100.0%

EPA Reg. No. 8033-141-70506

EPA SLN No. AZ-230302

ENVIRONMENTAL HAZARDS

This product is toxic to birds and aquatic invertebrates. This product is toxic to bees exposed to direct treatment. Do not apply this product while bees are foraging in the treatment area. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high-water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Do not contaminate water used for irrigation or domestic purposes.

THREATENED AND ENDANGERED SPECIES

Special local needs (SLN) labels are important to states like Arizona with niche industries and significant economic risks due to local pest problems. Our stewardship of this important labeling process is important and must also consider the potential off-target effects of any pesticide used in Arizona. This includes any potential harm to threatened or endangered (TE) species in our state. Please carefully consider the use of this pesticide and obtain as much information as you can to help ensure your application will have no impacts on TE species.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This supplemental label and the federally registered label must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on this supplemental label and the main EPA registered label.

USE	RATE (OZ PER ACRE)	PEST	COMMENTS
Cotton	4.5 – 6.7 oz/A (0.1-0.15 lb ai)	Whitefly	Begin applications when whitefly adults appear prior to development of nymphs. Do not wait until heavy populations have become established. For whitefly control, Assail 30SC Insecticide should be applied in a minimum finished spray volume of 5 gallons per acre by aircraft and 15 gallons per acre by ground equipment. Use the higher rate under heavy pressure. Make applications on a minimum 7 day interval as long as pest pressure continues. Whiteflies have shown a tendency to develop resistance. For resistance management purposes, alternating applications of different chemical classes reduces the potential for resistance development.

RESTRICTIONS AND PRECAUTIONS:

- Do not make more than 4 applications per calendar year.
- Do not apply more than once every 7 days.
- Do not apply less than 28 days before harvest (PHI = 28 days).
- Do not exceed a total of 0.4 lbs. a.i. (18.0 fl oz. of product) per acre per calendar year.
- There are no rotational crop plantback restrictions for this product.

Rev 3/29/2023

Sold By:
UPL NA Inc..
630 Freedom Business Center, Suite 402
King of Prussia, PA 19406
1-800-438-6071