

FIFRA Section 2(ee) Recommendation



FOR DISTRIBUTION AND USE ONLY IN THE STATES OF OREGON

BELAY[®] **INSECTICIDE**

EPA Reg. No. 59639-150

BELAY[®] INSECTICIDE FOR USE ON PACIFIC FLATHEADED BORER IN HAZELNUT (FILBERT) TREES

This FIFRA Section 2(ee) Recommendation Expires December 31, 2029.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This 2(ee) recommendation contains directions for use which do not appear on the package label and must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions and precautions on the EPA registered label when using *Belay* Insecticide alone or in tank mix combinations. The most restrictive labeling applies when using a tank mix.

***BELAY* INSECTICIDE APPLICATION TO HAZELNUT (FILBERT) TREES**

FOLIAR POST BLOOM APPLICATION		
Pest	<i>Belay</i> Insecticide Application Rate	Instructions
Pacific Flatheaded Borer	6 fl oz/A (0.1 lb ai/A)	Borer: Time application to the beginning of adult emergence. Apply <i>Belay</i> Insecticide to the entire tree including the trunk. Use adequate water volume and an adjuvant to help improve movement of <i>Belay</i> Insecticide into the bark.

RESTRICTIONS

- Do not apply more than 0.2 lb active ingredient clothianidin per acre per year.
- Do not apply treatments less than 10 days apart.
- Do not apply within 21 days of harvest.
- *Belay* Insecticide must not be applied during bloom or when bees are foraging.
- Do not feed or allow livestock to graze on cover crops from treated orchards.

Please contact Valent U.S.A. LLC at 800-6-VALENT (682-5368) to determine if this product is registered in your state.

©2019 Valent U.S.A. LLC

Registrant: Valent U.S.A. LLC
P.O. Box 5075
San Ramon, CA 94583

Belay is a registered trademark of Sumitomo Chemical Company, Ltd.