Product Bulletin

Corteva Agriscience LLC 9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

800-258-3033

Hemi SC®

EPA Reg. No.62719-545

2(ee) Recommendation[†]

For Distribution and Use Only in the State of TX

Suppression of Thrips in Cotton

This 2(ee) Recommendation expires December 31, 2027

ATTENTION

- † This recommendation is made as permitted under FIFRA 2(ee) and has not been submitted to or approved by the EPA.
- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- Read and follow all applicable directions for use, precautions, and limitations on the product label attached to the container for this product.

Directions for Use

Apply 1.5 to 3.0 fl oz of this product per acre to provide early season suppression of thrips in cotton. For heavier infestations, use a higher rate in the rate range.

Using Hemi SC is not a replacement for insecticide seed treatments or in-furrow insecticide applications at planting. Early season, foliar applications of Hemi SC to suppress thrips in cotton should only be made to fields that have already received an insecticide seed treatment or in-furrow insecticide application for thrips control.

Use of an adjuvant may increase the level of thrips control.

®™Trademarks of Corteva Agriscience and its affiliated companies

R260-003 Approved: 12/13/2024 **Initial Printing**