

## **Technical Information Bulletin**

Preplant applications to field corn with Distinct® herbicide

Use of a reduced rate of Distinct + glyphosate or 2,4-D ester in tank mixes for preplant burndown of emerged weeds.

FOR USE IN: AL, AR, CO, DE, GA, IA, IL, KS, KY, LA, MD, MI, MN, MO, MS, NE, NJ, NM, OH, OK, PA, SD, TN, VA, WI, WV, WY

EPA Reg. No. 7969-150

This recommendation is made as permitted under FIFRA section 2(ee) and had not been submitted to or approved by the EPA. All applicable directions, restrictions, precautions, and Conditions of Sale and Warranty on the EPA label are to be followed. Refer to the container label for additional instructions. Always read and follow label directions.

Information contained in this Technical Information Bulletin is not intended to replace or amend any product labeling. Always read and follow all label directions when using any pesticide alone or in mix combinations. The user must have this recommendation in their possession at the time of use.

## **DIRECTIONS FOR USE:**

**Distinct** may be applied preplant to field corn for control of emerged weeds. Do not apply on coarse soils with less than 2.5% organic matter.

Distinct 2 or 3 oz/acre + glyphosate*	Apply at least 14 days prior to planting	See main <b>Distinct</b> label for adjuvants required and weeds controlled**
--	--	--

<sup>\*</sup>See glyphosate label for application rate. Additional weeds controlled may be listed on the glyphosate label.

## OR

Distinct 2 or 3oz/acre + 2,4-D ester* (0.38-0.5	Apply at least 14 days prior to	See main <b>Distinct</b> label for adjuvants required and weeds controlled**
ae/acre)	planting	

<sup>\*</sup>See 2,4-D ester label for additional restrictions or limitations on corn planting intervals. Follow the most restrictive planting interval listed on the 2,4-D ester label or this **Distinct** 2(ee) recommendation.

© 2005 BASF Corporation All rights reserved Distinct is a registered trademark of BASF Corporation 007969-00150.20050216.**NVA 2005-04-078-0059** 

BASF Corporation Agricultural Products 26 Davis Drive Research Triangle Park, NC 27709



<sup>\*\*</sup>For glyphosate resistant marestail (*Conyza canadensis*), **Distinct** rate must be applied at 3 oz/acre for control of 4" (maximum height) marestail; for marestail taller than 4", **Distinct** rate must be 4 oz/acre

<sup>\*\*</sup>For improved control of common dandelion (Taraxacum officinale).