



## PRODUCT BULLETIN

### FIFRA SECTION 2(ee) RECOMMENDATION

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. Follow all applicable directions, restrictions, and precautions on the EPA-registered label. This product bulletin must be in the possession of the user at the time of application.

**Date:** May 30, 2025

**Product:** Phobos® FC EPA Reg. No. 89118-17 and AZterknot® Fungicide EPA Reg. No. 89118-14

**Use:** Phobos FC and AZterknot Fungicide tank mix for foliar disease control on soybean

**State:** AL, AR, DE, FL, GA, IL, IN, LA, MO, NC, NJ, OH, PA, SC, TN, VA

### DIRECTIONS FOR USE

#### SOYBEAN

**FOR BEST RESULTS, ONLY USE THE PHOBOS RATES LISTED BELOW WHEN PHOBOS FC IS TANKMIXED WITH AZTERKNOT FUNGICIDE (SEE RATE RANGE BELOW). ALTERNATIVELY, PHOBOS FC, AT THE RATES LISTED BELOW, CAN BE TANK-MIXED WITH AZTEROID FC FUNGICIDE, OTHER AZOXYSTROBIN FUNGICIDES OR OTHER GROUP 11 (STROBILURIN) FUNGICIDES REGISTERED FOR SOYBEANS FOR THE DISEASES LISTED BELOW.**

FOLIAR APPLICATION		
DISEASE	USE RATES	
Asian Soybean Rust ( <i>Phakopsora pachyrhizi</i> ) Frogeye Leaf Spot ( <i>Cercospora soja</i> ) Powdery Mildew ( <i>Microsphaera diffusa</i> ) Brown Spot ( <i>Septoria glycines</i> ) Alternaria Leaf Spot ( <i>Alternaria</i> species) Anthracnose ( <i>Colletotrichum truncatum</i> ) Cercospora Blight ( <i>Cercospora kikuchii</i> ) Pod and Stem Blight ( <i>Diaporthe phaseolorum</i> )	Phobos FC fl oz. product/acre (lb prothioconazole a.i. /A)	Azterknot fl oz. product/acre (lb azoxystrobin a.i./A   lb RSE/A)
	2.0 – 3.3 (0.046 – 0.078)	7.4 – 18.4 (0.1-0.25   0.06-0.14)



Rhizoctonia Aerial Blight  
(*Rhizoctonia solani*)

**Instructions:**

**Application Method:** PHOBOS FC + AZTERKNOT tank mix may be applied by ground, aerial or chemigation application equipment.

The lowest labeled rate of spray surfactant should be tank mixed with PHOBOS FC + AZTERKNOT to ensure optimal disease control.

**Application Timing:**

- For best results, apply the PHOBOS FC + AZTERKNOT tank mix as a broadcast foliar spray at early flowering or at first signs of disease development, whichever is earlier.

**RESTRICTIONS:**

- **AZTERKNOT:**
  - **Maximum Single Application:** **DO NOT** apply more than 18.4 fl oz/A (0.025 lb azoxystrobin and 0.14 lb RSE) per acre per application.
  - **Annual Maximum:**
    - **DO NOT** exceed 110.3 fl oz of AZterknot (1.5 lb azoxystrobin and 0.86 lb RSE) per acre per calendar year.
    - **DO NOT** exceed 1.5 lb azoxystrobin/A per calendar year from all azoxystrobin containing products.
    - **Forage and Hay:** **DO NOT** apply more than 1 application at 18.4 fl oz/A (0.25 lb azoxystrobin/A) per calendar year.
    - **DO NOT** apply more than 5 applications per calendar year at the high rate of 18.4 fl oz/A, or 14 applications per calendar year at the low rate of 7.4 fl oz/A.
- **PHOBOS FC:**
  - **Maximum Single Application:** **DO NOT** apply more than 6.66 fl oz of Phobos FC (0.156 lbs prothioconazole) per acre per foliar application.
  - **Annual Maximum**
    - **DO NOT** make more than 2 application per year.
    - **DO NOT** apply more than 7.6 fl oz of Phobos FC (0.178 lbs prothioconazole) per acre per application.
    - **DO NOT** apply more than 17.2 fl oz of Phobos FC (0.403 lbs prothioconazole) per acre per year.
- **DO NOT** make applications less than 14 days apart.
- **DO NOT** apply within 21 days of harvest
- **Resistance Management:** **DO NOT** make more than 2 sequential foliar application of any Group 11 fungicides before alternating to a fungicide with a different mode of action. **DO NOT** make more than 2 applications of any Group 11 fungicides per season.