



INDAZIFLAM

GROUP

29

HERBICIDE

FIFRA Section 24(c) Special Local Need Label**FOR DISTRIBUTION AND USE ONLY WITHIN GEORGIA****Application in citrus groves to trees established less than one year****Alion® Herbicide**

EPA Reg. No. 264-1106

SLN. No. GA-220003

**THIS LABEL IS VALID UNTIL DECEMBER 31, 2027 UNLESS OTHERWISE AMENDED,
WITHDRAWN, CANCELED, OR SUSPENDED.****ACTIVE INGREDIENT:**

Indaziflam*.....19.05%

OTHER INGREDIENTS:80.95%**TOTAL:**100.00%

Contains 1.67 pounds of indaziflam per gallon.

*(CAS No: 730979-19-8)

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This label must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

APPLICATION DIRECTIONS FOR USE IN CITRUS

Only apply Alion Herbicide in citrus groves where the soil has completely settled around citrus trees and there are no open channels or depressions in the soil that would allow the product to move into the root zone through open channels.

Citrus Crops: Crop group 10-10 including Australian desert lime; Australian finger-lime; Australian round lime; Brown River finger lime; calamondin; citron; citrus hybrids; clementine; grapefruit; Japanese summer grapefruit; kumquat; lemon; lime; Mediterranean mandarin; mount white lime; New Guinea wild lime; orange, sour; orange, sweet; pummelo; Russell River lime; satsuma mandarin; sweet lime; tachibana orange; Tahiti lime; tangelo; tangerine (mandarin); tangor; trifoliate orange; uniq fruit; cultivars, varieties, and or hybrids of these

Dose Rate Chart for Citrus

Soil Texture	Alion Herbicide (fl oz product / broadcast acre)
Any soil except those that contain 20% or greater gravel content	5.0 to 6.5 fl oz/A (0.065 to 0.085 lb ai/A)

USE RESTRICTIONS

- **Do not apply** more than 10.3 fl oz product/A (0.134 lb ai/A) per year or in a 12-month period.
- When making more than one application per year, allow a minimum of 90 days between applications.

USE PRECAUTIONS

- Avoid direct or indirect spray contact with crop foliage, green bark, roots, or fruit as it may cause localized crop injury or death. Only the trunks of trees transplanted more than one year may be sprayed with Alion Herbicide if the trunk is callused, mature brown bark. Contact of Alion Herbicide with tissues other than mature brown bark can result in serious damage or plant death.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577.

For **PRODUCT USE** Information Call 1-866-99BAYER (1-866-992-2937).

As with any crop-protection product, always read and follow label directions.

For additional information call toll-free 1-866-99BAYER (1-866-992-2937).



Bayer CropScience

Bayer CropScience LP
800 N. Lindbergh Blvd.
St. Louis, MO 63167
1-866-99BAYER (1-866-992-2937)