



## Section 2(ee) Recommendation

Date: March 2, 2011

Product: Renown® Fungicide, EPA Reg. No. 100-1315

Use: Renown on turfgrass for the preventive control of the additional diseases listed on this recommendation

State: AL, AR, AZ, CO, CT, DE, FL, GA, HI, IA, ID, IL, IN, KS, KY, LA, MA, ME, MD, MI, MO, MN, MS, MT, NC, ND, NE, NM, NJ, NV, NH, OH, OK, PA, RI, SC, SD, TN, UT, VA, VT, WA, WI, WV, WY

### Directions for Use

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- The user must refer to the federally approved labels for the above noted product and read and follow all directions for use, restrictions, and precautions.
- The user should have this recommendation in its possession at the time of use.
- Use of this product according to this bulletin has not been reviewed or endorsed by the Office of the Indiana State Chemist.
- This recommendation for use of this product is permitted under Section 2(ee) of FIFRA and has not been submitted to or been approved by EPA.

Renown provides preventive control of the following diseases for turfgrass species listed on the federal Section 3 label for Renown.

Target Diseases	Use Rate (fl oz product per 1000 sq ft)	Application Interval (Days)	Instructions
Brown Ring Patch ( <i>Waitea circinata</i> var. <i>circinata</i> )	2.5 – 4.5	14 - 28	Begin applications prior to disease development. For best control, use as a component of a preventive disease control program.
Fairy Ring ( <i>Lycoperdon</i> spp., <i>Agrocybe pediades</i> and <i>Bovistia plumbea</i> )	2.5 – 4.5	14 - 28	Apply as soon as possible after fairy ring symptoms develop. Apply only in 4 gallons water per 1000 square feet (174 gallons per acre). Add the recommended rate of a wetting agent to the final spray. Allow spray to dry, then irrigate with 0.25 inches of water prior to next mowing. Severely damaged or thin turf may require reseeding. Fairy ring symptoms may take 2 to 3 weeks to disappear following application.
Pythium Blight and Root Rot ( <i>Pythium aphanidermatum</i> and <i>Pythium</i> spp.)	4.5	10 - 14	Use preventatively. Begin applications before disease is present. During periods of prolonged favorable conditions, treat on the 10 day interval. For use on newly seeded as well as established turf.
Gray Snow Mold, Typhula Blight ( <i>Typhula incarnate</i> )	4.5	30	Apply prior to snow cover. Reapply at 30 day intervals if snow melts.
Pink Snow Mold ( <i>Microdochium nivale</i> )	4.5	30	Apply prior to snow cover. Reapply at 30 day intervals if snow melts.
Necrotic Ring Spot ( <i>Leptosphaeria korrae</i> )	4.5	30	Apply preventively in the Fall and Spring. Allow spray to dry, then irrigate with 0.25 inches of water prior to next mowing or within 24 hours of application.
Take-All Patch ( <i>Gaeumannomyces graminis</i> var. <i>avenae</i> )	4.5	30	Apply preventively in the Fall and Spring. Begin applications when conditions are favorable for disease infection. Allow spray to dry, then irrigate with 0.25 inches of water prior to next mowing or within 24 hours of application.
Summer Patch ( <i>Magnaporthe poae</i> )	4.5	30	Make 2-3 applications in the spring and summer. Initiate applications when soil temperatures reach 65°F at a 2-inch soil depth.

Renown® and the Syngenta logo are trademarks of a Syngenta Group Company.

©2011 Syngenta

2(ee) Registrant:  
Syngenta Crop Protection, LLC  
P. O. Box 18300  
Greensboro, NC 27419-8300

Label Code: AL1315077AC0311