

Fungicide/Bactericide

EPA Reg. No. 80289-3-10163

For the control of cryptogams on cranberry in the State of Wisconsin.

FIFRA §2(ee) RECOMMENDATION: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the United States Environmental Protection Agency (EPA).

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This product bulletin must be in the possession of the user at the time of pesticide application of Badge® SC Fungicide/Bactericide. This labeling contains supplemental instructions that may not appear on the product label. All applicable directions, restrictions, and precautions on the product label are to be followed.

CROP	DISEASE	RATE (PINTS PER ACRE)	MAX. APP. RATE	USE INSTRUCTIONS
Cranberry	Moss	3.5-7	44.4 pints product/acre 12.6 lbs Cu/acre for all disease applications	Apply postharvest and again in spring at bud swell. Apply 1 or 2 additional applications at 7 to 14 day intervals or longer as needed depending on disease severity. The minimum retreatment interval is 7 days. Where application rates and intervals are provided in a range (e.g. 8 to 24 pints and 7 to 10 days), the higher rates and shorter spray intervals are recommended when rainfall is heavy and/or disease pressure is high.

Badge is a registered trademark of Isagro ID# 19-11-01



Manufactured by Isagro SpA for: Isagro USA, Inc., 430 Davis Drive, Suite 240, Morrisville, NC 27560

Distributed by : Gowan Company LLC P.O. Box 5569 Yuma, AZ 85366-5569