

# SUPPLEMENTAL LABEL

## FOR USE ON BERRY AND SMALL FRUIT (SUBGROUP 13-07H) LOW GROWING BERRY SUBGROUP, EXCEPT STRAWBERRY



### KEEP OUT OF REACH OF CHILDREN CAUTION

- This labeling must be in the possession of the user at the time of pesticide application.
- It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.
- All applicable directions, restrictions, and precautions on the EPA-registered label are to be followed.
- 

#### DIRECTIONS FOR USE ON BERRY AND SMALL FRUIT (subgroup 13-07H) LOW GROWING BERRY SUBGROUP, EXCEPT STRAWBERRY

CROP	PEST	RATE	COMMENTS
Bearberry Bilberry Blueberry Lowbush Cloudberry Cranberry Lingonberry Muntries Partridgeberry Cultivars, varieties and/or hybrids of these	Blackheaded Fireworm (suppression only) Cranberry Fruitworm (suppression only) Cranberry Weevil (suppression only) Flea Beetles Leafhoppers Spanworm (suppression only) Sparganothis Fruitworm (suppression only) Stinkbugs Tipworm (suppression only)	<b>FOLIAR:</b> 3.5 to 7 fl oz/A (0.090 to 0.180 lb ai/A)	<b>Higher water volumes provide improved insect control.</b>  Begin applications when first pest activity is noticed or when insects reach threshold levels per State and County Extension Service recommendations. Repeat as needed to maintain control, but not more often than every 14 days. For best results, time application before a damaging population becomes established.  Under severe pest pressure, use the higher specified rates.  The rate applied affects the length of control. Use the high rate where infestations occur later in crop development, or where pest pressure is continuous. <i>SCORPION</i> can be mixed and/or alternated with commonly used insecticides, such as <i>Knack</i> , to improve length of control and/or achieve better knockdown of pests.
<b>NOTE: Foliar Application:</b> <ul style="list-style-type: none"> <li>• Apply with air or ground equipment in adequate water for uniform coverage (Use a minimum of 5 gals/A for air or 30 gals/A for ground applications).</li> <li>• Do not apply SCORPION within seven (7) days of harvest.</li> <li>• Do not apply more than a total of 14 fl oz/A of SCORPION (0.360 lb ai) per acre per season.</li> </ul>			

**Gowan**  
The Go To Company

EPA Reg. No. 10163-317  
EPA Est. No.

Distributed By:  
Gowan Company  
PO BOX 5569  
Yuma, AZ 85366