FUNGICIDE
TARIAN™

EPA Reg. No. 264-779-2935

For Distribution and Use Only Within the States of Illinois, Indiana, Iowa, Kansas, Kentucky, Michigan, Minnesota, Missouri, Nebraska, North Dakota, Oklahoma, Ohio, South Dakota, and Wisconsin

REDUCED APPLICATION RATE FOR EARLY SEASON CONTROL OF ANTHRACNOSE LEAF BLIGHT (*Colletotrichum graminicola*) AND EYESPOT (*Aureobasidium zeae*) IN CORN

ATTENTION
- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- Before using TARIAN™ in accordance with this recommendation, read and follow all applicable directions for use, restrictions, and precautions on the EPA registered label, as found on the package.
- This recommendation should be in the possession of the user at the time of application.
- This recommendation for use of this product is permitted under Section 2(ee) of FIFRA and has not been submitted to or been approved by EPA.

DIRECTIONS FOR USE

CORN (Field Corn, Popcorn, and Field Corn Grown for Seed)
For Anthracnose Leaf Blight (*Colletotrichum graminicola*) and Eyespot (*Aureobasidium zeae*) apply 5 – 10 fl oz per acre. Apply TARIAN as a broadcast foliar spray at V4 (four leaf collar) to V6 (6 leaf collar) growth stages when conditions are favorable for disease development. For season-long control of these and other diseases listed on the label, such as gray leaf spot and rust, apply a sequential treatment of this product at a rate of 10 fl oz per acre from growth stages VT through R2 (full tassel through blister). TARIAN may be applied by ground, air or chemigation.

Restrictions: Do not apply more than 24 fl oz of TARIAN per acre per season. Do not apply within 30 days of harvest for forage, grain and stover. Do not apply more than 2 sequential applications of TARIAN.

Use of this product according to this bulletin has not been reviewed or endorsed by the State of Indiana.

2023-0608

WILBUR-ELLIS®