



EPA Reg. No. 71512-31-88783

**Directions for Use on Crop Subgroup 13-07F (Grapes) to Control  
Black Rot and Phomopsis cane and leaf spot**

**PRODUCT BULLETIN**

**FIFRA Section 2(ee) Recommendation:** This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

**APPLICATION DIRECTIONS**

Crop	Diseases	Use Rate Fl. Oz. Product Per Acre	Instructions
Crop Subgroup 13-07F (Grapes)	Black rot ( <i>Guignardia bidwellii</i> )	8.5 fl oz	<b>Application Instructions:</b> For use on all types of grapes (wine, table, raisin, and juice). Apply when important for disease control, ideally prior to disease initiation. Refer to university or extension publications for ideal timings. Apply with sufficient water to allow for penetration into the foliage to obtain complete coverage using 50 to 100 gallons of spray volume per acre. Continue as needed on a 14 to 21 day interval. When disease pressure is severe, use shorter intervals.  <b>Resistance Management:</b> Do not make more than 2 sequential applications of this product or other Group 3 or Group 7 containing fungicides before rotating to a fungicide with a different mode of action. Do not apply a third application of this product within 28 days of the second application.  <b>Restrictions:</b> Do Not apply more than 25.5 fl oz/A/year The Pre-Harvest Interval (PHI) for this crop is 14 days. This product may be applied via aerial application equipment except in the State of New York. Do Not apply more than the maximum yearly rate from any combination of products containing Isofetamid (1.72 lb. a.i. per acre per year) or Tebuconazole (0.9 lb. a.i. per acre per year) to these crops.
	Phomopsis cane and leaf spot ( <i>Phomopsis viticola</i> )	8.5 fl oz	

Includes all members of the Fruit, Small Vine Climbing Crop Subgroup 13-07F, except fuzzy kiwifruit: Amur river grape; gooseberry; grape; kiwifruit, hardy; maypop; schisandra berry; and cultivars, varieties, and/or hybrids of these.

For **24-Hour Medical Emergency Assistance** (Human or Animal) Call 1-888-484-7546.  
 For **Chemical Emergency**, Spill, Leak, Fire or Accident, Call CHEMTREC 1-800-424-9300.

Read and follow all applicable directions for use, precautions and limitations on the product label attached to the FERVENT 475SC Fungicide container.



Summit Agro USA, LLC  
 240 Leigh Farm Road, Suite 215  
 Durham, NC 27707

Fervent is an invention and registered trademark of Ishihara Sangyo Kaisha, Ltd., and is manufactured and developed by ISK Biosciences Corporation.