



TECHNICAL INFORMATION BULLETIN  
FOR THE USE OF MAJESTENE® BIONEMATICIDE  
ON PINEAPPLE

**FIFRA Section 2(ee) Recommendation**

This recommendation is made as permitted by Section 2(ee) of FIFRA, as amended, and has not been submitted to or approved by the U. S. Environmental Protection Agency.

FOR USE IN ALL STATES WHERE CROP AND PEST(S) EXIST AND WHERE 2(EE) RECOMMENDATIONS ARE RECOGNIZED.

**MAJESTENE®**  
**EPA Reg. No. 84059-14**

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

| Crop      | Target Nematodes  | Application Method                                      | Product Use Rate per Application       | Application Instructions  |
|-----------|---|---|--|---|
| Pineapple | Suppression of plant parasitic nematodes including root lesion, root knot, ring, stubby, sting and dagger nematodes | In-furrow or banded over an open furrow during planting | 1 - 2 gallons per acre in-furrow<br>or | Apply Majestene at a rate of 1-2 gallons per acre in a directed or banded spray over an open furrow. Consult the Majestene label for proper rates for each row width.   |
|           |   | Soil drench during transplant                           | 1-2% v/v drench<br>or                  | Drench transplants with a 1-2% v/v dilution of Majestene during transplanting with sufficient water to thoroughly soak the root zone.   |
|           |   | In-season directed spray during cultivation             | 2 gallons per acre<br>or               | Apply Majestene at a rate of 2 gallons per acre in a spray directed to the base of the plant during cultivation operations with spray directed in front of cultivator shovels to insure incorporation into the soil. Consult the Majestene label for proper rates for each row width. |
|           |   | In-season chemigation                                   | 2 gallons per acre                     | Apply Majestene at a rate of 2 gallons per acre through overhead irrigation   |

MAJESTENE® has a **pre-harvest interval (PHI)** of **0 days**.

Do not enter or allow worker entry into treated areas during the **restricted-entry interval (REI)** of **4 hours**.



**Nematode management considerations** - To obtain maximum results in a nematode management program, it is important to take soil samples prior to application to assess nematode populations and determination of an appropriate management strategy. Utilization of the services of a field person or crop consultant who is knowledgeable in nematode management is advised.

**All applicable directions, restrictions and precautions on the EPA-registered label are to be followed.**

This FIFRA Section 2(ee) recommendation contains new or additional directions for use of this product, which may not appear on the package label. Read and carefully observe the cautionary statements plus all the other information appearing on the product label.

**This recommendation must be in the possession of the user at the time of pesticide application.**

MAJESTENE® is a registered trademark of Marrone Bio Innovations, Inc.

© Copyright Marrone Bio Innovations, Inc., 2014

Marrone Bio Innovations, Inc.

1540 Drew Ave.

Davis, California 95618

Telephone (530) 750-2800

[www.marronebioinnovations.com](http://www.marronebioinnovations.com)