



# PRODUCT BULLETIN

## FIFRA 2(ee) Recommendation

### FOR USE AND DISTRIBUTION IN THE STATE OF CALIFORNIA ONLY

**August 28, 2018**  
**OxiDate 2.0**  
**EPA Reg. No. 70299-12**

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or accepted by the U.S. Environmental Protection Agency. This product bulletin must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

#### Use on Herbs & Spices to Control:

**Anthrachnose, Downy Mildew, Powdery Mildew, Pythium Rot**

#### Directions For Use:

*It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. This product bulletin contains new or additional directions for use which are recommended by BioSafe Systems, LLC. under FIFRA Section 2(ee).*

Crop	Disease	Dilution Rate	Application Notes
<b>Herbs and Spices</b> Basil Chives Cilantro Coriander Dill Mint Rosemary Sage	Anthrachnose Downy Mildew Powdery Mildew Pythium Rot	1:500 to 1:1,000	<b>Preventive:</b> For best results, apply at first sign of disease. Use 12.8 fl. oz. to 25.6 fl. oz of OxiDate 2.0 per 100 gallons of water. <b>For ground applied application.</b> Apply 30-100 gallons of spray solution per acre. Ensure adequate coverage of upper and lower foliage, stems, branches and stalks. Maintain a 5-day interval spray cycle until harvest. <b>For aerial applied applications:</b> Apply 3-20 gallons of spray solution per acre. Do not make applications at a height greater than 10 ft. above the plant canopy, unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to wind and evaporation. Do not exceed the maximum application rate or apply more often than labeled in the Application Instructions for that crop.