

Product Bulletin



Corteva Agriscience LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Pixxaro® EC

EPA Reg. No. 62719-735

Special 2(ee) Recommendation†

FOR USE AND DISTRIBUTION ONLY IN THE STATE OF OREGON

Crop Rotation Intervals

Refer to product label for Pixxaro® EC for general use precautions, mixing directions, application directions, and restrictions.

ATTENTION

†This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- Read the label affixed to the container for Pixxaro EC insecticide before applying. Carefully follow all precautionary statements and applicable use directions except as specified below.
- Use of Pixxaro EC according to this Section 2(ee) recommendation is subject to all use precautions and limitations imposed by the label affixed to the container for Pixxaro EC.

The following rotational crops may be planted at the indicated interval following application of this product. For best results conduct a field bioassay prior to planting any broadleaf crops not listed. **DO NOT** plant unlisted crops prior to 15 months prior to application.

Crop	Rotation Interval ¹ (Months)
barley, triticale, wheat (spring, winter, and durum)	0
field corn, sweet corn	3 days
grass grown for seed, forage or hay, oats, sorghum	14 days
alfalfa ² , canola, cotton, millet, popcorn, rice, rye, seed corn, soybean, sugarcane, sunflower	4
brassica (cole) leafy vegetables, camelina, chickpea, clover, dry bean, flax, mustard, peanut, peas (dry and succulent), safflower, sugar beet	9
Potato (not for seed) ³	10
other crops not listed	15

¹Minimum number of months that must pass before planting other crops after application of Pixxaro EC.

² For rotation to alfalfa, cumulative precipitation (including irrigation) must be greater than 6.0 inches between application date and alfalfa seeding date. Otherwise, rotation to alfalfa is recommended 9 months following application.

³For rotation to potatoes, precipitation (including irrigation) must be greater than 8.0 inches during the 10 months following application of Pixxaro EC. Otherwise, rotation to potatoes is recommended 15 months following application.

™®Trademarks of Corteva Agriscience and its affiliated companies

R445-005

Issued: 03/04/24

Initial Printing