



UPL NA, Inc.
630 Freedom Business Center
Suite 402
King of Prussia, PA 19406

FIFRA Section 2(ee) Recommendation

VACCIPLANT®

Reduced Rate in Corn (Field, Pop, Seed, Silage, Sweet), Soybeans, Wheat, Barley, and Rice EPA Reg. No. 70506-608

For distribution and use only within the states of Alabama, Alaska, Arizona, Arkansas, Colorado, Connecticut, Delaware, Florida, Georgia, Idaho, Illinois, Indiana, Iowa, Kansas, Kentucky, Louisiana, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New Mexico, North Carolina, North Dakota, Ohio, Oklahoma, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This recommendation is made as permitted by FIFRA Section 2(ee) and has not been submitted to or approved by the U.S. Environmental Protection Agency.

This 2(ee) recommendation, contains directions for use, which do not appear on the package label and must be in the possession of the user at the time of VACCIPLANT® application.

Follow all applicable directions, restrictions, Worker Protection Standard requirements and precautions on the EPA registered label.

Use of this product according to this bulletin has not been reviewed or endorsed by the Office of the Indiana Chemist.

VACCIPLANT can be used at a reduced rate of 7 fl oz/A in corn (field, pop, seed, silage, sweet), Soybeans, Wheat, Barley, and Rice when tank-mixed with another registered fungicide.

APPLICATION INSTRUCTIONS

Crop	Application Method	Product Use Rate per Application/Acre	Application Instructions
Corn Sweet Corn Field Corn Popcorn Silage Corn Seed Corn Soybean Wheat Barley Rice	Foliar	7 fl oz	To optimize disease control and maximize yields, apply VACCIPLANT, at different stages of crop. For example, in Corn, prior to tassel, In Rice, at late boot to pinnacle emergence. Start VACCIPLANT applications preventatively or when conditions for disease development become favorable. Reapply every 7 – 14 days. Under moderate to heavy disease pressure, tank mix this product with another registered fungicide.
<ul style="list-style-type: none">Follow all applicable directions, restrictions, and precautions on the labels of all products used in the tank-mix. This includes mixing instructions.VACCIPLANT has a pre-harvest interval (PHI) of 0 days.Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 hours.			

UPL, the UPL logo and VACCIPLANT are trademarks of a UPL Corporation Limited Group Company.