## RESTRICTED USE PESTICIDE

**Due to Toxicity to Fish and Aquatic Organisms** 

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.



Bayer CropScience LP P.O. Box 12014 2 T.W. Alexander Drive Research Triangle Park, North Carolina 27709 1-866-99BAYER (1-866-992-2937) GROUP 3 INSECTICIDE

Baythroid® XL EPA Reg. No. 264-840

For control of Brown Marmorated Stink Bug (BMSB) in Grape and Potato.

## PRODUCT BULLETIN

# FOR DISTRIBUTION AND USE ONLY IN THE STATES OF INDIANA, MICHIGAN, MINNESOTA, OHIO, AND WISCONSIN

**FIFRA Section 2(ee) Recommendation:** This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or accepted by the U.S. Environmental Protection Agency. This product bulletin should be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

#### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product inconsistent with its labeling.

GRAPE Includes: Table grape, Raisin, Wine and Muscadin	e grape	
PEST CONTROLLED	Rate fluid ounces/Acre	Rate Ib Al/Acre
Brown Marmorated Stink Bug	2.4 – 3.2	0.019 - 0.025

## Foliar Application Restrictions

Pre-Harvest Interval (PHI): 3 days.

Maximum BAYTHROID XL allowed per 14-day interval: 3.2 fluid oz/A (0.025 lb Al/Acre). Maximum BAYTHROID XL allowed per crop season: 12.8 fluid oz/A (0.1 lb Al/Acre). Minimum application volume (water): 50 GPA – ground, 10 GPA – aerial application.

#### POTATO AND OTHER TUBEROUS AND CORM VEGETABLES

## Includes all members of Crop Subgroup 1C:

Arracacha, Arrowroot, Artichoke (Chinese and Jerusalem), Edible canna, Cassava (bitter and sweet), Chayote root, Chufa, Dasheen (taro), Ginger, Leren, Potato, Sweet potato, Tanier, True yam, Turmeric, Yam bean

(tare), enger, ceren, retate, ewest petate, ramer, rrae jam, ramene, ram sear		
PEST CONTROLLED	Rate	Rate
	fluid ounces/Acre	lb Al/Acre
Brown Marmorated Stink Bug	1.6 – 2.8	0.013 - 0.022

### Foliar Application Restrictions

Pre-Harvest Interval (PHI): 0 day.

If more than 5.6 fluid oz/Acre is applied, allow at least 14 days between last application and grazing.

Maximum BAYTHROID XL allowed per 5-day interval: 2.8 fluid oz/A (0.022 lb Al/Acre).

Maximum BAYTHROID XL allowed per crop season: 16.8 fluid oz/A (0.132 lb Al/Acre).

For <u>MEDICAL</u> And <u>TRANSPORTATION</u> Emergencies <u>ONLY</u> Call 24 Hours A Day 1-800-334-7577 For <u>PRODUCT USE</u> Information Call 1-866-99BAYER (1-866-992-2937)

As with any crop-protection product, always read and follow label directions. For additional information call toll-free 1-866-99BAYER (1-866-992-2937)