



Bayer CropScience LP
P.O. Box 12014
2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
1-866-99BAYER (1-866-992-2937)

Stratego® YLD Fungicide

Reduced Application Rate for Early Season
Control of Anthracnose Leaf Blight
(*Colletotrichum graminicola*) and
Eyespot (*Aureobasidium zeae*) in Corn

EPA Reg. No. 264-1093

PRODUCT BULLETIN

**FOR DISTRIBUTION AND USE ONLY WITHIN THE STATES OF ARKANSAS,
COLORADO, DELAWARE, ILLINOIS, INDIANA, IOWA, KANSAS, KENTUCKY,
LOUISIANA, MARYLAND, MICHIGAN, MINNESOTA, MISSOURI, MONTANA,
NEBRASKA, NEW JERSEY, NEW YORK, NORTH CAROLINA, NORTH DAKOTA,
OHIO, OKLAHOMA, PENNSYLVANIA, SOUTH CAROLINA, SOUTH DAKOTA,
VIRGINIA, WISCONSIN AND WYOMING**

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. It is a violation of federal law to use this product in a manner inconsistent with its label. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

DIRECTIONS FOR USE

CORN (Field Corn, Popcorn, and Field Corn Grown for Seed)	
Disease Control	Rate fluid ounces/acre
Anthracnose Leaf Blight (<i>Colletotrichum graminicola</i>)	2.0 – 5.0
Eyespot (<i>Aureobasidium zeae</i>)	

Notes and Restrictions: Apply Stratego YLD Fungicide as a broadcast foliar spray at V4 (four leaf collar) to V7 (7 leaf collar) growth stages when conditions are favorable for disease development. For season-long control of these and other diseases listed on the label, such as gray leaf spot and rust, apply a sequential treatment of Stratego YLD Fungicide at 4 to 5 fluid ounces per acre from growth stages VT through R2 (full tassel through blister).

Stratego YLD Fungicide may be applied by ground, air or chemigation.

Do not apply more than 10 fluid ounces of Stratego YLD Fungicide per acre per season. Do not apply within 14 days of harvest for grain and fodder. Forage may be harvested the same day of application.

Do not apply more than 2 sequential applications of Stratego YLD Fungicide.

Use of this product according to this bulletin has not been reviewed or endorsed by the State of Indiana.

For **MEDICAL** And **TRANSPORTATION** Emergencies **ONLY** Call 24 Hours A Day 1-800-334-7577
For **PRODUCT USE** Information Call 1-866-99BAYER (1-866-992-2937)

As with any crop-protection product, always read and follow label directions.
For additional information call toll-free 1-866-99BAYER (1-866-992-2937).