

# FIFRA Section 24(c) Special Local Need Label

Corteva Agriscience LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

For Distribution and Use Only Within the State of California  
For Use on Cherry for the Control of Spotted Wing Drosophila

## Entrust<sup>®</sup> SC

EPA Reg. No. 62719-621

SLN CA-130001

### ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This state-specific Section 24(c) labeling must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, and precautions on the Entrust SC label (EPA Reg. No 62719-621) and this label.

### Directions for Use

**Application Timing:** Treat when pests appear, targeting eggs at hatch or small larvae. Consult your company representative, extension service specialist, certified crop advisor or your state agricultural experiment station for any additional local use recommendations for your area.

**Application Rate:** Apply 4.8 to 6.4 fl. oz of Entrust SC per acre as a foliar spray to control spotted wing drosophila. Use a higher rate in the rate range for larger larvae or moderate to severe pest infestations. Heavy infestations may require repeat applications but follow resistance management guidelines.

**Resistance Management:** Do not make more than 2 consecutive foliar applications of Group 5 insecticides (spinosad and spinetoram). If additional applications are required, rotate to an effective product in a different group for at least 2 applications. Do not make more than three foliar applications per acre per year of Group 5 insecticides (spinetoram [Delegate<sup>®</sup>] and spinosad [Entrust<sup>®</sup>, Success<sup>®</sup>]).

**Treatment Schedule:** If three Entrust SC applications are made, the first may be made up to 28 days before harvest, the second up to 10 days before harvest, and the final application at 3 days before harvest. Two applications of one or more effective insecticides not in Group 5 must be made between the first and second Entrust SC applications.

### Specific Use Restrictions

- **Pre-Harvest Interval:** Do not apply within 3 days of harvest.
- Do not apply more than 6.4 fl oz of Entrust SC (0.10 lb ai) per acre per application.
- Do not apply more than a total of 12.8 fl oz of Entrust SC (0.20 lb ai) per acre within 10 days of harvest.
- Do not apply more than a total of 19.2 fl oz of Entrust SC (0.30 lb ai) per acre per year.
- Do not make more than three foliar applications per acre per year of Group 5 insecticides (spinetoram and spinosad). This restriction supersedes the number of applications on the label affixed to the container of Entrust SC.
- Apply by air blast sprayer. Do not apply by hand wand or backpack sprayer.

<sup>™</sup><sup>®</sup>Trademarks of Corteva Agriscience and its affiliated companies

Corteva Agriscience LLC  
9330 Zionsville Road  
Indianapolis, IN 46268  
1-800-992-5994

R399-049  
Accepted: 05/24/2023  
Replaces R399-034