

Product Bulletin

Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Intrepid Edge®

EPA Reg. No. 62719-666

(For Use in California)

Special 2(ee) Recommendation†

Dormant or Delayed-Dormant applications of Intrepid Edge for the Control of Peach Twig Borer (PTB) in Almonds

ATTENTION

†This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- Read the label affixed to the container for Intrepid Edge® insecticide before applying. Carefully follow all precautionary statements and applicable use directions except as specified below.
- Use of Intrepid Edge according to this Section 2(ee) recommendation is subject to all use precautions and limitations imposed by the label affixed to the container for Intrepid Edge.

Application Timing: Apply Intrepid Edge as a dormant or delayed-dormant foliar spray targeting PTB in the hibernacula or when they have exited the hibernacula, but prior to penetrating shoots. Lepidopterous larvae must be controlled before they penetrate the shoots.

Begin dormant or delayed-dormant applications after January 1st. Consult your Corteva Agriscience representative, extension service specialist, certified crop advisor or your state agricultural experiment station for any additional local use recommendations for your area.

Use of Crop Oils: Crop oils labeled for agricultural use may be added to the dormant spray solution for suppression of overwintering mites and scale insects. Consult specific oil labels and University or Extension recommendations for precautions and restrictions regarding the use of oils in nut and fruit trees.

Application Rate: The amount of Intrepid Edge applied per acre will depend upon tree size, volume of foliage present and pest pressure. Use a higher rate in the rate range for larger trees or heavy infestations.

Pest	Intrepid Edge® (fl oz/acre)
Peach twig borer	6 – 12

Restrictions:

- **Preharvest Interval:** Do not apply within 7 days of harvest.
- Do not apply more than 18 fl oz of Intrepid Edge® per acre per application.
- Do not apply more than a total of 51 fl oz of Intrepid Edge® (1 lb ai methoxyfenozide, 0.2 lb ai spinetoram) per acre per season.
- **Minimum Treatment Interval:** Do not make applications less than 14 days apart.
- **Maximum Number of Applications:** Do not make more than 4 applications per year.
- Do not make more than two consecutive applications of Group 5 or Group 18 insecticides within a crop season.

®TM Trademarks of Dow AgroSciences, DuPont or Pioneer and their affiliated companies or respective owners

R911-012

Issued: 01/14/2020