



Supplemental Label

BIFENTURE® EC Agricultural Insecticide

(EPA Reg. No. 70506-57)

DIRECTIONS FOR USE ON TOBACCO

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Follow all applicable directions, restrictions and precautions on the federally-registered label.

This label must be in the possession of the user at the time of pesticide application.

PESTS CONTROLLED	RATE BIFENTURE EC/ACRE	APPLICATION INSTRUCTIONS	PHI
Armyworm spp. Cutworm spp. Mole Crickets Stalkborers Tobacco Flea Beetle (larvae) White Grubs Wireworms	0.0625-0.10 lb ai (4.0-6.4 fl. oz. product)	Pre-transplant soil application: Apply specified rate in a minimum of 10 gal/A. To control below-ground pests, use equipment which will incorporate the application into the top 4" of soil. At-transplant water treatment application: Apply specified rate in a water treatment volume of 10-200 gal/A.	Do not apply later than layby. Do not apply more than 0.2 lb ai/A per season.
Aphid spp. Armyworm spp. Chinch Bugs Cucumber Beetles Flea Beetle (Adults) Stink Bugs Japanese Beetles Grasshoppers Cutworm spp. Tarnished Plant Bugs Green Bugs Saltmarsh Caterpillar Thrips Tobacco Budworm* Tobacco Hornworm Whiteflies	0.04-0.10 lb ai (2.56-6.4 fl. oz. product)	Foliar application: Apply specified rate in a minimum of 10 gal/A up to and including layby. Do not make more than 2 foliar applications per season. May be tank mixed with other herbicides registered for tobacco use. *See Resistance section on this label for more information.	
Lygus spp. Spider Mites	0.1 lb ai (6.4 fl. oz. product)		

Rev. 5/09
CA appr. 10/28/09

United Phosphorus, Inc.
630 Freedom Business Center
King of Prussia, PA 19406
1-800-438-6071 • www.upi-usa.com