

Recommendation

FIFRA SECTION 2(ee) RECOMMENDATION

GROUP 28 INSECTICIDE

Shenzi 700 WG Insecticide

(EPA Reg. No. 70506-609)

FOR SUPPRESSION OF CARPOPHILUS BEETLE Carpophilus truncatus IN TREE NUTS IN THE STATE OF CALIFORNIA

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by USEPA.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read and follow all applicable instructions, restrictions, and precautions, and Conditions of Sale and Warranty on the EPA-approved label. This label and the federally registered label must be in the possession of the user at the time of pesticide application.

CROP	APPLICATION METHOD	INSECT	SHENZI 700 WG INSECTICIDE RATE PER ACRE IN FL OZ (LB A.I.)	RE- TREATMENT INTERVAL IN DAYS (RTI)	PRE- HARVEST INTERVAL IN DAYS (PHI)	RE-ENTRY INTERVAL IN HOURS (REI)
Tree Nuts, (EPA Crop Group 14- 12)	Foliar	Carpophilus Beetle	2.2 (0.098)	7	10	4

APPLICATION INSTRUCTIONS:

Spray Volume:

- Thorough coverage is essential. Select a spray volume appropriate for the size of trees or plants and density of foliage.
- DO NOT apply less than 30 gals water per acre.
- For best results apply 100 150 gals water per acre by ground. Where higher spray volumes are used, apply a higher SHENZI WG INSECTICIDE rate in the specified rate range.

Carpophilus Beetle (larva) (Hull-split application timing): Make an application at 1 - 5% hull-split timing; make a second application approximately 10 - 14 days later. Depending on level of pest infestation, use of higher rates in the labelled rate range and multiple applications may be needed

Grazing on Tree Nut Orchard or Grove Floor: There are no grazing restrictions for (1) Grass forage, fodder, and hay. Any grass Gramineae family (either green or cured) except sugarcane and those included in the cereal grains group, that will be fed to or grazed by livestock, all pasture and range grasses and grasses grown for hay or silage, and (2) Nongrass animal feeds.

RESTRICTIONS:

- DO NOT make more than 4 applications per acre per calendar year.
- DO NOT apply more than 4.6 oz of SHENZI WG INSECTICIDE or 0.2 lb a.i. chlorantraniliprole containing products per acre per calendar year.