

Product Bulletin



Corteva Agriscience LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Rezuvant[®]

EPA Reg. No. 62719-750

Special 2(ee) Recommendation[†]

Crop Rotation Intervals

(For Use In: AK, AL, AR, AZ, CO, DE, GA, IA, ID, IL, IN, KS, KY, LA, MD, MI, MN, MO, MS, MT, NC, ND, NE, NJ, NM, NV, OH, OK, OR, PA, SC, SD, TN, UT, VA, WA, WI, WV, WY)

Refer to product label for Rezuvant[®] for general use precautions, mixing directions, application directions, and restrictions.

ATTENTION

[†]This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- Read the label affixed to the container for Rezuvant insecticide before applying. Carefully follow all precautionary statements and applicable use directions except as specified below.
- Use of Rezuvant according to this Section 2(ee) recommendation is subject to all use precautions and limitations imposed by the label affixed to the container for Rezuvant.

The following rotational crops may be planted at the indicated interval following application of this product. For best results conduct a field bioassay prior to planting any broadleaf crops not listed. Do not plant unlisted crops less than 15 months prior to application.

Crop	Rotation Interval ⁽¹⁾ (Months)
barley, wheat (spring, winter, and durum)	0
field corn, sweet corn, triticale, grass grown for seed, forage or hay, oats, sorghum	3
alfalfa ² , canola, cotton, millet, popcorn, rice, rye, seed corn, soybean, sugarcane, sunflower	4
brassica (cole) leafy vegetables, camelina, chickpea, clover, dry bean, flax, mustard, peanut, peas (dry and succulent), safflower, sugar beet	9
Potato (not for seed) ³	10
other crops not listed	15

¹Minimum number of months that must pass before planting other crops after application of Rezuvant.

²For rotation to alfalfa, cumulative precipitation (including irrigation) must be greater than 6.0 inches between application date and alfalfa seeding date. Otherwise, rotation to alfalfa is recommended 9 months following application.

³For rotation to potatoes, precipitation (including irrigation) must be greater than 8.0 inches during the 10 months following application of Rezuvant. Otherwise, rotate to potatoes a minimum of 15 months following application.

[™][®]Trademarks of Corteva Agriscience and its affiliated companies

R472-001
Issued: 06/06/2024
Initial Printing