



Section 2(ee) Recommendation

Date: March 6, 2024

Product: Velista®
EPA Reg. No. 100-1534

Use: Addition of Diseases as Pests Controlled on Turfgrasses

State: AK, AL, AR, AZ, CA, CO, CT, DC, DE, FL, GA, HI, IA, ID, IL, IN, KS, KY, LA, MA, MD, ME, MI, MN, MO, MS, MT, NC, ND, NE, NH, NJ, NM, NV, OH, OK, PA, RI, SC, SD, TN, UT, VA, VT, WA, WI, WV, WY

Directions for Use

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- The user must refer to the federally approved labels for the above noted product and read and follow all directions for use, restrictions, and precautions.
- Use of this product according to this bulletin has not been reviewed or endorsed by the Office of the Indiana State Chemist. The user should have this recommendation in its possession at the time of use.
- This recommendation for use of this product is permitted under Section 2(ee) of FIFRA and has not been submitted to or been approved by EPA.

Disease	Application Site	Rate oz/1,000 sq ft	Application Interval (Days)	Comments
Rapid Blight (<i>Labyrinthula terrestris</i>)	All Turfgrass	0.5	14	Apply when conditions are favorable for disease development. Do not apply more than 2.2 oz product/1,000 sq ft/year.
Fairy Ring (<i>Basidiomycetes</i>)	All Turfgrass	0.5-0.7	14-28	Apply as part of a preventative program. Rotate with Heritage® and Briskway® fungicides. Do not apply more than 2.2 oz product/1,000 sq ft/year.
Leaf and Sheath Spot (<i>Rhizoctonia zeae</i>)	All Turfgrass	0.3-0.5	14	Use as part of a preventative program in tank mixture and/or rotation with Daconil Action®, Heritage and Briskway fungicides. Do not apply more than 2.2 oz product/1,000 sq ft/year.

Disease	Application Site	Rate oz/1,000 sq ft	Application Interval (Days)	Comments
Spring Dead Spot (<i>Ophiosphaerella</i> spp.)	All Turfgrass	0.5-0.7	28	Make two applications on a 28 day interval prior to winter dormancy. Do not apply more than 2.2 oz of product/1,000 sq ft/year.
SummerPatch (<i>Magnaporthe poae</i>)	All Turfgrasses	0.3-0.5	14-28	During periods of high disease pressure, tank mix with a fungicide that is registered for the control of summer patch. Apply when conditions are favorable for disease development. Do not apply more than 2.2 oz of product/1,000 sq ft/year.
Brown Ring Patch (<i>Waitea circinata</i> var <i>circinata</i>)	All Turfgrasses	0.5	14	Apply when conditions are favorable for disease development. Do not apply more than 2.2 oz of product/1,000 sq ft/year.
Large Patch (<i>Rhizoctonia solani</i>)	All Turfgrasses	0.5-0.7	14-28	Make 1-2 applications in the fall as a preventative. Apply an early spring application if disease pressure is high. Do not apply more than 2.2 oz of product/1,000 sq ft/year.

Velista®, Briskway®, Daconil Action®, Heritage® and the Syngenta logo are trademarks of a Syngenta Group Company.

©2024 Syngenta

2(ee) Registrant:
 Syngenta Crop Protection, LLC
 P. O. Box 18300
 Greensboro, NC 27419-8300

Label Code: AK1534011BC0324