

RESTRICTED USE PESTICIDE

DUE TO TOXICITY TO NON-TARGET INVERTEBRATES, MAMMALS, AND AQUATIC ORGANISMS

FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION, AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION.

syngenta.

ITERA

Section 24(c) Special Local Need Label

ABAMECTIN	GROUP	6	INSECTICIDE
-----------	-------	---	-------------

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF COLORADO

Agri-Mek® SC Miticide/Insecticide

For Control of Spider Mites (including Banks Grass Mite) Infesting Grasses Grown for Seed Only

**EPA Reg. No. 100-1351
EPA SLN No. CO-190001**

This label expires and must not be distributed or used in accordance with this SLN registration after December 31, 2029

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This label must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA-registered label.

IMPORTANT

No portion of the crop treated with Agri-Mek SC Miticide/Insecticide shall be used or distributed as food for human consumption or feed for animal consumption.

Syngenta has not investigated the use of Agri-Mek SC Miticide/Insecticide for potential adverse interactions with any other crop protection or fertilizer products used in grass seed production, nor across potential experimental or commercial breeding lines. Therefore adverse effects arising from the use of Agri-Mek SC on grasses grown for seed and cannot be predicted and are therefore the responsibility of the User.

GRASSES GROWN FOR SEED

Note:

- Apply Agri-Mek SC with ground application equipment or aircraft
- For best control apply Agri-Mek SC with ground application equipment. With aerial application the resulting control of spider mites could be less than with ground application.
- Agricultural products formulated with binder or sticker type components may reduce performance of Agri-Mek SC.

Pest	Rate per Acre per Application	Instructions
Spider Mites (Tetranychid species including Banks Grass Mite)	2.5 - 3.5 fl. oz./A	<ul style="list-style-type: none">• Agri-Mek SC must be mixed with a non-ionic activator type wetting, spreading and/or penetrating spray adjuvant. Do not use binder or sticker type adjuvants. (See Use Information section found at the beginning of the Agri-Mek SC label and Use Restrictions below.)• Apply when spider mites are first observed.• Repeat applications as needed to maintain control within the constraints of a sound resistance management program

USE RESTRICTIONS

- **DO NOT** cut current year's treated seed crop for hay or forage.
- **DO NOT** allow livestock to graze in treated areas.
- **DO NOT** graze or feed livestock on cover crops growing in treated areas.
- **DO NOT** allow Agri-Mek SC treated screenings from seed processing to enter food or feed channels.
- **DO NOT** allow Agri-Mek SC treated seed for sprouting.
- **All seed treated with Agri-Mek SC is to be tagged** at processing plants "NOT FOR HUMAN OR ANIMAL CONSUMPTION". It shall be the responsibility of the grower to inform the processing plants of any seed treated with Agri-Mek SC.
- **Pre-Harvest Interval (PHI):** Do not apply within **21 days** of harvest for seed.
- **Adjuvant Requirement:** Agri-Mek SC **must** be mixed with a spray adjuvant as instructed in the **Use Information** section found at the beginning of the Agri-Mek SC label. The spray adjuvant must be approved for use in grass seed production. Carefully follow the **Directions for Use** and **Precautions** on the spray adjuvant label as well as the precautions in the official local spray guides. Do not use binder or sticker type adjuvants.
- **Application Method:** Ground or aerial application is permitted.
- **Amount of Water:** Use sufficient water volume to ensure good coverage of foliage. Do apply in less than 20 gallons of water/A with ground application equipment. Do apply in less than 10 gallons of water/A with aircraft. Under conditions such as high pest populations, dense foliage, or adverse weather conditions (such as high temperatures), use a greater volume of water to ensure adequate coverage.

- **Maximum Amount per Application:** Do not apply more than 3.5 fl. oz./A (or 0.019 lb. a.i./A) of Agri-Mek SC or any other foliarly applied abamectin-containing product per application.
 - **Application Interval:** Wait at least 7 days before repeating application.
 - **Number of Applications:** For resistance management, do not make more than 2 sequential applications of Agri-Mek SC or any other foliarly applied abamectin-containing product.
 - **Maximum Amount per Season:** Do not exceed a total of 7 fl. oz./Acre of Agri-Mek SC or 0.038 lb ai of abamectin per acre in a growing season.
-

©2024 Syngenta

Agri-Mek® and the Syngenta logo are trademarks of a Syngenta Group Company

24(c) Registrant:
Syngenta Crop Protection, LLC
P. O. Box 18300
Greensboro, NC 27419-8300

Label Code: CO1351126BA1224