



EPA REG. NO. 66222-257

FIFRA §2(ee) RECOMMENDATION

Strawberries Use Rate for Botrytis and Leaf Spot
FIFRA Section 2(ee) recommendation for use in the following state(s): FLORIDA

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This recommendation must be in the possession of the user at the time of pesticide application.
- Read the "Limitation of Warranty and Liability" in the label booklet for Captan Gold 4L before using.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by U.S. EPA.

STRAWBERRIES		
DISEASES	APPLICATION RATE CAPTAN GOLD 4L QUARTS/ ACRE	APPLICATION INSTRUCTIONS
Botrytis Leaf Spot	1.5 – 3.0	<p>Ground Application - Apply by broadcast in sufficient water for thorough coverage.</p> <p>Air application – Apply in 10 to 20 gallons of water.</p> <p>Begin applications when new growth starts in the spring and before fruit starts to form. Repeat at 7 to 14 day intervals.</p> <p>Under conditions favorable to fruit rot, continue applications up to day of harvest period treating immediately after each picking.</p>
<p>If applying as a directed/banded spray; use band rate of Captan GOLD 4L according to the following formula:</p> $\frac{\text{Plant Bed Width (inches)}}{\text{Row Spacing (inches)}} \times \text{Broadcast rate per acre} = \text{Banded rate of Captan GOLD 4L per acre}$		
<p>RESTRICTIONS:</p> <ul style="list-style-type: none"> • REI: 24 hours • PHI: 0 day. May be applied up to day of harvest. • Do not apply more than 24 quarts per Acre per crop cycle 		

103114