

THIS RECOMMENDATION IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE EPA.

2(ee) Recommendation



FOR DISTRIBUTION AND USE ONLY IN
Oregon, Washington, Idaho, Montana, Utah

EPA REG. NO. 84876-2

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS
PRODUCT IN A MANNER INCONSISTENT WITH ITS
LABELING

This Product Bulletin Contains New or Additional
Directions For Use Which Are Recommended By
Sym-Agro, Inc. Under FIFRA Section 2(ee)

CROP	PEST	RATE OF APPLICATION
Pome Fruit (Crop Group 11)	Gray mold (Botrytis cinerea)	40 - 45.7 fl oz/A (0.66-0.75 lb ai/A)
DIRECTIONS FOR USE: Pre-Harvest use to Control Post Harvest Disease: Apply as a pre-harvest spray within 10 to 1 days of harvest. Thorough coverage of the fruit is required. Application closer to harvest may provide better efficacy. General: Increase spray volume as growth increases to provide thorough coverage of foliage and fruit for optimum disease control.		
RESTRICTIONS: Do not make more than two sequential applications of ProBlad® Verde Fungicide before alternating to a labeled fungicide with a different mode of action. Do not make more than 5 applications of ProBlad® Verde Fungicide per harvest cycle.		
PHI: 1 Day		
ALL APPLICABLE DIRECTIONS, RESTRICTIONS, WORKER PROTECTION STANDARDS REQUIREMENTS AND PRECAUTIONS ON THE EPA REGISTERED PROBLAD VERDE® FUNGICIDE (EPA REG. NO. 84876-2) LABEL MUST BE FOLLOWED. THIS RECOMMENDATION MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.		



Sym-Agro, Inc.
PO Box 71
Visalia, CA 93291

ProBlad is a trademark of CEV S.A.

ProBlad Verde_2ee_Pome Fruit Pre-Harvest_07-25-2020