

FIFRA Section 2(ee) Label

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the U.S. Environmental Protection Agency

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF FLORIDA

CARBARYL 4L EPA REG NO. 34704-447

For control of Asian Citrus Psyllid in Citrus

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Crop	Pest	Amount of Carbaryl 4L per acre	Specific Directions
Citrus (grapefruit, lemons, limes, oranges, tangelos, tangerines, citrus citron, kumquat, and hybrids)	Asian Citrus psyllid (<i>Diaphorina</i> citri)	1.5 - 3.0 qts/A	Repeat applications as necessary up to a total of 8 times but not more often than once every 14 days. Use the lower rate on light to moderate infestations, young plants and early instars, and use the higher rate on heavy infestations, mature plants, advanced instars and adults. Thorough and uniform spray coverage is essential for effective control.

Use Restrictions and Precautions:

- Do not apply within 5 days of harvest.
- Do not apply more than a total of 20 quarts per acre per crop.
- For ground application, apply in sufficient volume for adequate coverage.
- For aerial application, use at least 10 gallons of spray mixture per acre for adequate distribution.
- Observe Bee Caution.

Loveland Products, Inc P.O. Box 1286 Greeley, CO 80632